



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification:</b> <b>A61K 41/00</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 00/62808</b> <b>(43) International Publication Date:</b> 26 October 2000 (26.10.2000)
<b>(21) International Application Number:</b> PCT/US00/10100 <b>(22) International Filing Date:</b> 15 April 2000 (15.04.2000) <b>(30) Priority Data:</b> 60/129,733 16 April 1999 (16.04.1999) US 60/159,873 15 October 1999 (15.10.1999) US <b>(60) Parent Application or Grant</b> MAYO OF FOUNDATION FOR MEDICAL EDUCATION AND RESEARCH [/]; (). REGENTS OF THE UNIVERSITY OF MINNESOTA [/]; (). COLLINS, Douglas, A. [/]; (). HOGENKAMP, Henricus, P., C. [/]; (). COLLINS, Douglas, A. [/]; (). HOGENKAMP, Henricus, P., C. [/]; (). VIKSNINS, Ann, S.; ().		<b>Published</b>
<b>(54) Title: COBALAMIN CONJUGATES USEFUL AS ANTITUMOR AGENTS</b> <b>(54) Titre: CONJUGUES DE COBALAMINE UTILISES COMME AGENTS ANTITUMORAUX</b>  <b>(57) Abstract</b> <p>The invention provides cobalamin compounds linked to a neutron capture therapy target (e.g. Boron-10 or Gadolinium-157), and optionally linked to a detectable moiety, as well as pharmaceutical compositions comprising the compounds, and methods for using the compounds in medical diagnosis and therapy.</p> <b>(57) Abrégé</b> <p>L'invention concerne des composés de cobalamine liés à une cible thérapeutique de capture de neutrons (Bore-10 ou Gadolinium-157), et éventuellement liés à une fraction décelable, ainsi que des compositions pharmaceutiques comprenant lesdits composés, et des méthodes d'utilisation de ces composés dans le diagnostic et les thérapies médicales.</p>		

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau


## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>7</sup> : <b>A61K 41/00</b>		(11) International Publication Number: <b>WO 00/62808</b>
<b>A2</b>		(43) International Publication Date: <b>26 October 2000 (26.10.00)</b>
(21) International Application Number: <b>PCT/US00/10100</b>		(71)(72) Applicants and Inventors: <b>COLLINS, Douglas, A.</b> [US/US]; 1150 Meadowlark Court S.W., Rochester, MN 55902 (US). <b>HOGENKAMP, Henricus, P., C.</b> [US/US]; 2211 Marion Road, Roseville, MN 55113 (US).
(22) International Filing Date: <b>15 April 2000 (15.04.00)</b>		
(30) Priority Data:		
60/129,733	16 April 1999 (16.04.99) US	
60/159,873	15 October 1999 (15.10.99) US	
(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Applications		(74) Agent: <b>VIKSINNS, Ann, S.</b> ; Schwegman, Lundberg, Woessner & Kluth, P.O. Box 2938, Minneapolis, MN 55402 (US).
US	60/129,733 (CIP)	
Filed on	16 April 1999 (16.04.99)	
US	60/159,873 (CIP)	
Filed on	15 October 1999 (15.10.99)	
(71) Applicants (for all designated States except US): <b>MAYO OF FOUNDATION FOR MEDICAL EDUCATION AND RESEARCH</b> [US/US]; 200 First Street S.W., Rochester, MN 55905 (US). <b>REGENTS OF THE UNIVERSITY OF MINNESOTA</b> [US/US]; 600 University Gateway, 200 Oak Street SE, Minneapolis, MN 55455-2070 (US).		(81) Designated States: <b>AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO</b> patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), <b>Eurasian</b> patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), <b>European</b> patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), <b>OAPI</b> patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
		Published Without international search report and to be republished upon receipt of that report.

(54) Title: **COBALAMIN CONJUGATES USEFUL AS ANTITUMOR AGENTS**

## (57) Abstract

The invention provides cobalamin compounds linked to a neutron capture therapy target (e.g. Boron-10 or Gadolinium-157), and optionally linked to a detectable moiety, as well as pharmaceutical compositions comprising the compounds, and methods for using the compounds in medical diagnosis and therapy.



**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LJ	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## D scription

5

10

15

20

25

30

35

40

45

50

55

## COBALAMIN CONJUGATES USEFUL AS ANTITUMOR AGENTS

**Related Application**

This application claims priority to U.S. Provisional Application Ser. No. 60/129,733, filed 16 April 1999; and U.S. Provisional Application Ser. No. 60/159,873, filed 15 October 1999.

**Background of the Invention**

Boron neutron capture therapy is based on the nuclear reaction that occurs when a stable isotope,  $^{10}\text{B}$ , is irradiated with low energy (0.025 eV) or thermal neutrons to yield helium nuclei ( $\alpha$ -particles) and  $^7\text{Li}$  nuclei.

The therapeutic potential of this reaction was recognized by Locher over 50 years ago (Locher, G.L. et al., Am. J. Roentgenol. Radium Ther., 36, 1-13 (1936)), but it was Sweet (Javid, M. et al., J. Clin. Invest., 31, 603-610 (1952); Sweet, W.H., N. Engl. J. Med., 245, 875-878 (1951); Sweet, W.H. et al., J. Neurosurg., 2, 200-209 (1952)), who first suggested that boron neutron capture therapy (BNCT) might be useful for the treatment of brain tumors.

Shortly thereafter, a clinical trial was initiated at the Brookhaven National Laboratory in cooperation with Sweet and others at the Massachusetts General Hospital utilizing borax as the capture agent (Farr, L.E. et al., Am. J. Roentgenol., 71, 279-291 (1954); Godwin, J.T. et al., Cancer (Phila.), 8, 601-615 (1955)). The objective at that time was to use BNCT as an adjunct to surgery for the treatment of patients with the most highly malignant and therapeutically refractory of all brain tumors, glioblastoma multiforme.

Further trials were carried out in the early 1960s, but these failed to show any evidence of therapeutic efficacy (Farr, L.E. et al., *supra*; Godwin, J.T. et al., *supra*; Asbury, A.K. et al., J. Neuropathol. Exp. Neurol., 31, 278-303 (1972)) and were associated with adverse effects in normal tissues (Asbury, A.K. et al., *supra*). Stimulated by the more encouraging clinical studies of Hatanaka et al. (Hatanaka, H.A., J. Neurol., 202, 81-94 (1975); Hatanaka, H. et al., Boron Neutron Capture Therapy for Tumors, Chap. 25, pp. 349-378. Niigata, Japan:

5 Nishimura Co., Ltd (1986)) for the treatment of malignant gliomas and the use of  
Mishima et al. (Mishima, Y. et al., *Lancet*, 2, 388-289 (1989)) for melanoma,  
there has been renewed national and international interest in BNCT.

10 The theoretical advantage of BNCT is that it is a two component  
or binary system, consisting of  $^{10}\text{B}$  and thermal neutrons, which when combined  
together generate high linear energy transfer (LET) radiation capable of  
selectively destroying tumor cells without significant damage to normal tissues.  
15 In order for BNCT to succeed a critical amount of  $^{10}\text{B}$  and a sufficient number of  
thermal neutrons must be delivered to individual tumor cells.

20 Over the past few years the Department of Energy and the NIH  
have renewed funding for BNCT-related research, and this has supported a  
growing number of investigators in many different disciplines. Advances in  
BNCT in the areas of compound distribution and pharmacokinetics compare  
favorably with other emerging modalities such as photon activation therapy,  
15 photodynamic therapy, and the use of radiolabeled antibodies for cancer  
treatment in which physiological targeting is used.

25 There are a number of nuclides that have a high propensity for  
absorbing low energy or thermal neutrons, and this property, referred to as the  
neutron capture cross-section ( $\sigma$ ), is measured in barns ( $1 \text{ b} = 10^{-24} \text{ cm}^2$ ). Of the  
30 various nuclides that have high neutron capture cross-sections,  $^{10}\text{B}$  is the most  
attractive for the following reasons: (a) it is nonradioactive and readily  
available, comprising approximately 20% of naturally occurring boron: (b) the  
particles emitted by the capture reaction [ $^{10}\text{B}(n,\alpha)^7\text{Li}$ ] are largely high LET: (c)  
35 their path lengths are approximately 1 cell diameter (10-14  $\mu\text{m}$ ), theoretically  
limiting the radiation effect to those tumor cells that have taken up a sufficient  
amount of  $^{10}\text{B}$  and simultaneously sparing normal cells and (d) the extensive  
chemistry of boron is such that it can be incorporated into a multitude of  
40 different chemical structures.

45  $^7\text{Li}$  and  $\alpha$ -particles are the primary fission product of the neutron  
capture reaction with  $^{10}\text{B}$ .  $\alpha$ -Particles are relatively slow and give rise to closely  
spaced ionizing events that consist of tracks of sharply defined columns. They  
have a path length of approximately 10  $\mu\text{m}$ , are high LET, and destroy a wide  
50 variety of biologically active molecules including DNA, RNA, and proteins. For

5 these reasons there is little, if any, cellular repair from  $\alpha$ -particle-induced radiation injury.

Since the  $^{10}\text{B}(n,\alpha)^7\text{Li}$  reaction will produce a significant biological effect only when there is a sufficient fluence of thermal neutrons and a critical amount of  $^{10}\text{B}$  localized around, on, or within the cell, the radiation produced can be extremely localized thereby sparing normal tissue components. Thus, selectivity is simultaneously one of the advantages and disadvantages of BNCT, since it requires delivery of boron-10 to tumor cells in greater amounts than normal cells.

10 Ideally, boron compounds to be used for BNCT should have a high specificity for malignant cells with concomitantly low concentrations in adjacent normal tissues and blood. Since it is desirable to confine the radiation solely to these cells, an intracellular and optimally intranuclear localization of boron would be preferred.

15 Several boron-containing derivatives of chlorpromazine have been synthesized (Nakagawa, T. et al., Chem. -Pharm. Bull. (Tokyo), 24, 778-781 (1976); Alam, F. et al., Sthralenther. Onkol., 165, 121-125 (1989)) and are being evaluated for their *in vivo* tumor localizing properties. *p*-Boronophenylalanine is another compound that is being studied as a potential capture agent for the treatment of melanoma. The rationale for its use is the avidity of melanomas for aromatic amino acids and their subsequent incorporation into melanin (Ichihashi, M. et al., J. Invest. Dermatol., 78, 215-218 (1982); Mishima, Y. et al., Neutral Capture Therapy, 230-236, Niigata, Japan: Nishimura Co., Ltd. (1986)).

25 Tumor localization has been demonstrated following i.v. administration by means of whole body autoradiography (Coderre, J.A. et al., Cancer Res., 48, 6313-6316 (1988)) and in several patients with cutaneous melanoma following perilesional injection (Mishima, Y. et al., Sthralenther. Onkol., 165, 251-254 (1989)). Stimulated by Mishima's experience, a number of other boron-containing amino acids have been synthesized that potentially could be incorporated in larger amounts into proteins of malignant cells (Hall, I.H. et al., J. Pharm. Sci., 68, 685-688 (1979)).

Another approach to the selective targeting of boron to melanomas is based on the observation that thiouracil is preferentially incorporated into melanotic melanomas during melanogenesis (Whittacker, J.R., J. Biol. Chem., 246, 6217-6226 (1971)). This observation provided the impetus for the synthesis of several boron-containing thiouracils (Gabel, D., Clinical Aspects of Neutron Capture Therapy, 233-241, New York: Plenum Publishing Co. (1989)), and these currently are being evaluated in animals.

Two other classes of compounds with a propensity for localizing in malignant tumors are the porphyrins and the related phthalocyanines. The biochemical basis by which these compounds achieve elevated concentration in malignant tumors is unknown, but this observation has served as the rationale for the use of hematoporphyrin derivative in the photodynamic therapy of cancer (Dougherty, T.J. et al., Porphyrin Photosensitization, 3-13, New York: Plenum Publishing Corp. (1981)).

The high concentration of these compounds in tumors and their intracellular localization and persistence have stimulated several groups of investigators to synthesize boronated porphyrins (Kahl, S. B. et al., Neutron Capture Therapy, 61-67, Niigata, Japan: Nishimura Co., Ltd. (1986)) and phthalocyanines (Alam, F. et al., Strahlenther. Oncol., 165, 121-123 (1989)) as potential capture agents. Boronated porphyrins appear to be 3-4 times more effective per unit dose in cell culture than the monomeric or dimeric form of  $\text{Na}_2\text{B}_{12}\text{H}_{11}\text{SH}$  (Laster, B.H. et al., Strahlenther. Oncol., 165, 203-205 (1989)). Although liver concentrations of these compounds are also high (Kahl, S.B. et al., *supra*) this would not limit their use as a capture agent for the treatment of brain tumors.

One final category of low molecular weight boron compounds are boron-containing purines and pyrimidines and their nucleosides. The rationale for their development is that such compounds may be selectively incorporated into rapidly proliferating tumor cells and trapped within the cell following their conversion to the corresponding nucleotide. Alternatively, these bases and their nucleosides may function as analogues of naturally occurring precursors of nucleic acids and become incorporated into nuclear DNA.



5                   Cytoplasmic or preferably a nuclear localization of all of these  
boron compounds would be advantageous since the heavy particles resulting  
from the capture reaction would deliver a greater proportion of their energy to  
10                   intranuclear targets, thereby permitting lower boron concentrations than would  
5                   have been required if the compounds were located extracellularly (Gabel, D. et  
al., Radiat. Res., 111, 14-25 (1987); Fairchild, R.G. et al., Int. J. Radiat. Oncol.  
Biol. Phys., 11, 831-840 (1985)). Schinazi and Prusoff (Schinazi, R.F. et al.,  
15                   Tetrahedron Lett., 50, 4981-4984 (1978)) have synthesized the first boron-  
containing nucleoside, 5-dihydroxyboryl-2'-deoxyuridine, an analogue of  
10                   thymidine, and have shown that it was not cytotoxic to African green monkey  
(Vero) cells at a concentration level of 1600  $\mu$ M (Laster, B.H. et al., Neutron  
Capture Therapy, 46-54, Niigata, Japan: Nishimura Co., Ltd. (1986)). *In vitro*  
20                   neutron radiation studies of cells grown in the presence of 5-dihydroxy-2'-  
deoxyuridine produced a biological effect that was equivalent to a concentration  
15                   of  $\sim 6 \mu\text{g } ^{10}\text{B/g}$ , which, if attainable *in vivo*, would be sufficient for BNCT.

25                   During the 1960s and early 1970s, interests developed in the  
potential use of polyclonal antibodies directed against tumor-associated antigens  
for the delivery of drugs and radioisotopes to tumors (Pressman, D. et al., Cancer  
Res., 40, 3001-3007 (1957); Ghose, T. et al., Br. Med. J., 1, 90-93 (1967);  
30                   Ghose, T. et al., Cancer (Phila.), 29, 1398-1400 (1972)). In 1964, Soloway  
suggested that antibodies might be used for the selective targeting of  $^{10}\text{B}$  to  
tumors (Soloway, A.H., *supra*). Hawthorne et al. (Hawthorne, M.F. et al., J.  
Medicinal Chem., 15, 449-452 (1972)) reported on the incorporation of the  
35                   diazonium salt from 1-(4-aminophenyl)-1,2-dicarbo-*closo*-dodecaborane into  
25                   antibodies directed against bovine serum albumin and polyclonal antibodies  
directed against human and mouse histocompatibility antigens (Hawthorne,  
40                   M.F. et al., *supra*).

                  It was claimed from *in vitro* experiments that these  
immunoconjugates were capable of delivering enough boron to human and  
45                   30                   murine lymphocytes to sustain a lethal  $^{10}\text{B}(n,\alpha)^7\text{Li}$  reaction, as evidenced by  
reduced viability following neutron irradiation. However, the  
immunoconjugates contained only 0.2% natural boron by weight, which was  
50                   equal to 6 atoms of  $^{10}\text{B}$ /molecule of antibody. In retrospect, it appears that there

5 must have been some other explanation for the reduced cell viability that was  
observed. Sneath et al. (Sneath, R.L., Jr., J. Medicinal Chem., 17, 796-799  
(1974)) showed that water-solubilizing groups had to be incorporated into  
10 protein-binding polyhedral boranes if protein solubility in aqueous systems was  
5 to be maintained.

Subsequently, a group of polyhedral borane derivatives  
containing protein-binding functional groups were linked to IgG molecules by  
means of the carbodiimide reaction without evidence of precipitation (Sneath,  
15 R.L. et al., J. Medicinal Chem., 19, 1290-1294 (1976)).

10 One final category of macromolecular species that are possibly  
useful for the delivery of  $^{10}\text{B}$  is what may be termed "encapsulating complexes,"  
such as liposomes, microspheres, and low density lipoproteins (Kahl, S.B. et al.,  
20 Strahlenther. Onkol., 165, 137-139 (1989)). Theoretically, large amounts of  $^{10}\text{B}$   
could be encapsulated, and if these encapsulating complexes could be targeted to  
15 the tumor by linkage to a monoclonal antibody using existing methodology or  
targeting an endogenously expressed cell surface receptor, they might be  
25 powerful delivery systems. Again, there may be preferential localization in the  
reticuloendothelial system, and methodology would have to be developed to  
30 minimize this and maximize tumor uptake.

#### 20 Cobalamin

For several years after the isolation of vitamin  $\text{B}_{12}$  as  
cyanocobalamin in 1948, it was assumed that cyanocobalamin and possibly  
35 hydroxocobalamin, its photolytic breakdown product, occurred in man. Since  
then it has been recognized that cyanocobalamin is an artifact of the isolation of  
25 vitamin  $\text{B}_{12}$  and that hydroxocobalamin and the two coenzyme forms,  
methylcobalamin and adenosylcobalamin, are the naturally occurring forms of  
40 the vitamin.

The structure of these various forms is shown in Figure 1,  
wherein X is CN, OH,  $\text{CH}_3$ , or adenosyl, respectively. Hereinafter, the term  
45 30 cobalamin will be used to refer to all of the molecule except the X group. The  
fundamental ring system without cobalt (Co) or side chains is called *corrin* and  
the octadehydrocorrin is called *corrole*. Figure 1 is adapted from The Merck  
50 Index, Merck & Co. (11th ed. 1989), wherein X is above the plane defined by the

5 corrin ring and nucleotide is below the plane of the ring. The corrin ring has attached six amidoalkyl ( $H_2NC(O)Alk$ ) substituents, at the 2, 3, 7, 8, 13, and 18 positions, which can be designated a-e and g, respectively. See D.L. Anton et al., I. Amer. Chem. Soc., 102, 2215 (1980).

10 5 Methylcobalamin serves as the cytoplasmic coenzyme for  $^3N$ -methyltetrahydrofolate:homocysteine methyl transferase (methionine synthetase, EC 2.1.1.13), which catalyzes the formation of methionine from homocysteine. Adenosylcobalamin is the mitochondrial coenzyme for methylmalonyl CoA mutase (EC 5.4.99.2) which interconverts methylmalonyl CoA and succinyl  
15 CoA.

20 All forms of vitamin  $B_{12}$  (adenosyl-, cyano-, hydroxo-, or methylcobalamin) must be bound by the transport proteins, Intrinsic Factor and Transcobalamin II to be biologically active. Specifically, gastrointestinal absorption of vitamin  $B_{12}$  relies upon the intrinsic factor-vitamin  $B_{12}$  complex  
25 being bound by the intrinsic factor receptors in the terminal ileum. Likewise, intravascular transport and subsequent cellular uptake of vitamin  $B_{12}$  throughout the body is dependent upon transcobalamin II and the cell membrane transcobalamin II receptors, respectively. After the transcobalamin II-vitamin  
30  $B_{12}$  complex has been internalized, the transport protein undergoes lysozymal degradation, which releases vitamin  $B_{12}$  into the cytoplasm. All forms of vitamin  $B_{12}$  can then be interconverted into adenosyl-, hydroxo-, or methylcobalamin depending upon cellular demand. See, for example, A.E. Finkler et al., Arch. Biochem. Biophys., 120, 79 (1967); C. Hall et al., J. Cell Physiol., 133, 187 (1987); M.E. Rappazzo et al., J. Clin. Invest., 51, 1915 (1972) and R. Soda et al.,  
35 Blood, 65, 795 (1985).

40 Cells undergoing rapid proliferation have been shown to have increased uptake of thymidine and methionine. (See, for example, M.E. van Eijkeren et al., Acta Oncologica, 31, 539 (1992); K. Kobota et al., J. Nucl. Med., 32, 2118 (1991) and K. Higashi et al., J. Nucl. Med., 34, 773 (1993)). Since  
45 methylcobalamin is directly involved with methionine synthesis and indirectly involved in the synthesis of thymidylate and DNA, it is not surprising that methylcobalamin as well as Cobalt-57-cyanocobalamin have also been shown to have increased uptake in rapidly dividing tissue (for example, see, B.A. Cooper  
50

et al., *Nature*, 191, 393 (1961); H. Flodh, *Acta Radiol. Suppl.*, 284, 55 (1968); L. Bloomquist et al., *Experientia*, 25, 294 (1969)). Additionally, upregulation in the number of transcobalamin II receptors has been demonstrated in several malignant cell lines during their accelerated thymidine incorporation and DNA synthesis (see, J. Lindemans et al., *Exp. Cell Res.*, 184, 449 (1989); T. Amagasaki et al., *Blood*, 26, 138 (1990) and J.A. Begly et al., *J. Cell Physiol.*, 156, 43 (1993).

Vitamin B<sub>12</sub> has several characteristics which potentially make it an attractive *in vivo* tumor therapeutic agent. Vitamin B<sub>12</sub> is water soluble, has no known toxicity, and in excess is excreted by glomerular filtration. In addition, the uptake of vitamin B<sub>12</sub> could potentially be manipulated by the administration of nitrous oxide and other pharmacological agents (D. Swanson et al., *Pharmaceuticals in Medical Imaging*, MacMillan Pub. Co., NY (1990) at pages 621-628).

A process for preparing <sup>125</sup>I-vitamin B<sub>12</sub> derivatives is described in U.S. Pat. No. 3,981,863 issued to Niswender et al. In this process, vitamin B<sub>12</sub> is first subjected to mild hydrolysis to form a mixture of monocarboxylic acids, which Houts, *infra*, disclosed to contain mostly the (e)-isomer. The mixture is then reacted with a p-(aminoalkyl)phenol to introduce a phenol group into the B<sub>12</sub> acids (via reaction with one of the free carboxylic acid groups). The mixed substituent B<sub>12</sub> derivatives are then iodinated in the phenol-group substituent. This U.S. patent teaches that the mixed <sup>125</sup>I-B<sub>12</sub> derivatives so made are useful in the radioimmunoassay of B<sub>12</sub>, using antibodies raised against the mixture.

U.S. Pat. No. 4,465,775 issued to T. M. Houts reported that the components of the radiolabelled mixture of Niswender et al. did not bind with equal affinity to IF. Houts disclosed that radioiodinated derivatives of the pure monocarboxylic (d)-isomer are useful in assays of B<sub>12</sub> in which IF is used. However, although Houts generally discloses that the monocarboxylic (d)-isomer can be labelled with fluorophores or enzymes and used in competitive assays for vitamin B<sub>12</sub> in fluids, a continuing need exists for labelled vitamin B<sub>12</sub> derivatives suitable for tumor and organ imaging and therapy.

U.S. Patent Number 5,739,313 issued to Collins and Hogenkamp reported that cobalamin analogs comprising a linking group and a chelating

group optionally comprising a detectable radionuclide or a paramagnetic ion  
localize in tumor cells and are useful for imaging tumors.

Despite previous efforts to identify a neutron capture agent that  
localizes in tumor cells in high concentration and is useful to treat cancer, there is  
currently a need for neutron capture agents that are useful to treat tumors.

#### Summary of the Invention

The invention provides a compound of the invention which is a  
residue of a compound of formula I (Figure 1) I linked to a residue of a molecule  
comprising Boron-10 (i.e., B-10), wherein X is CN, OH, CH<sub>3</sub>, adenosyl, or a  
molecule comprising B-10; or a pharmaceutically acceptable salt thereof.

The invention also provides a compound wherein a residue of a  
compound of formula I (Figure 1) is linked to a group of the formula Q-L-W-  
Det; wherein X is CN, OH, CH<sub>3</sub>, adenosyl, a molecule comprising B-10, or Q-L-  
W-Det; wherein Det is a chelating group comprising Gd-157; L is a linker or  
absent; and W and Q are each independently -N(R)C(=O)-, -C(=O)N(R)-,  
-OC(=O)-, -C(=O)O-, -O-, -S-, -S(O)-, -S(O)<sub>2</sub>-, -C(=O)-, -N(R)-, or a direct  
bond; wherein each R is independently H or (C<sub>1</sub>-C<sub>6</sub>)alkyl; or a pharmaceutically  
acceptable salt thereof.

The invention also provides a compound wherein a residue of a  
compound of formula I (Figure 1) is linked to a residue of a molecule comprising  
B-10; wherein a residue of the compound of formula I is also linked to a group  
of the formula Q-L-W-Det, wherein X is CN, OH, CH<sub>3</sub>, adenosyl, a group of the  
formula Q-L-W-Det, or a molecule comprising B-10; wherein: Det is a chelating  
group comprising a therapeutic radionuclide or a diagnostic radionuclide; L is a  
linker or absent; and Q and W are each independently -N(R)C(=O)-,  
-C(=O)N(R)-, -OC(=O)-, -C(=O)O-, -O-, -S-, -S(O)-, -S(O)<sub>2</sub>-, -C(=O)-, -N(R)-,  
or a direct bond; wherein each R is independently H or (C<sub>1</sub>-C<sub>6</sub>)alkyl; or a  
pharmaceutically acceptable salt thereof.

The invention also provides a compound wherein a residue of a  
compound of formula I (Figure 1) is linked to a residue of a molecule comprising  
B-10; wherein the residue of the compound of formula I is also linked to a group  
of the formula Q-L-W-Det, wherein X is CN, OH, CH<sub>3</sub>, adenosyl, a group of the  
formula Q-L-W-Det, or a molecule comprising B-10; Det is a chelating group

comprising Gd-157; L is a linker or absent; and Q and W are each independently -N(R)C(=O)-, -C(=O)N(R)-, -OC(=O)-, -C(=O)O-, -O-, -S-, -S(O)-, -S(O)<sub>2</sub>-, -C(=O)-, -N(R)-, or a direct bond; wherein each R is independently H or (C<sub>1</sub>-C<sub>6</sub>)alkyl; or a pharmaceutically acceptable salt thereof.

The invention also provides a compound wherein a residue of a compound of formula I (Figure 1) is linked to a detectable radionuclide and also linked to a residue of a molecule comprising Boron-10 (i.e., B-10), wherein X is CN, OH, CH<sub>3</sub>, adenosyl, or a molecule comprising B-10; or a pharmaceutically acceptable salt thereof.

The invention also provides a compound wherein a residue of a compound of formula I (Figure 1) is linked to a detectable radionuclide; and is also linked to a group comprising Gd-157; wherein X is CN, OH, CH<sub>3</sub>, adenosyl, a molecule comprising B-10, or a group comprising Gd-157; or a pharmaceutically acceptable salt thereof. Specifically, the group comprising Gd-157 can have the formula Q-L-W-Det; wherein Det is a chelating group comprising Gd-157; L is a linker or absent; and W and Q are each independently -N(R)C(=O)-, -C(=O)N(R)-, -OC(=O)-, -C(=O)O-, -O-, -S-, -S(O)-, -S(O)<sub>2</sub>-, -C(=O)-, -N(R)-, or a direct bond; wherein each R is independently H or (C<sub>1</sub>-C<sub>6</sub>)alkyl.

The invention also provides a compound wherein a residue of a compound of formula I (Figure 1) is linked 1) to a molecule comprising B-10 or to a chelating group comprising Gd-157, and is linked 2) to at least one residue of the formula Q-L-W-Det; wherein each Det is independently a chelating group comprising a metallic radionuclide; each L is independently a linker or absent; and each W and Q is independently -N(R)C(=O)-, -C(=O)N(R)-, -OC(=O)-, -C(=O)O-, -O-, -S-, -S(O)-, -S(O)<sub>2</sub>-, -C(=O)-, -N(R)-, or a direct bond; wherein each R is independently H or (C<sub>1</sub>-C<sub>6</sub>)alkyl; or a pharmaceutically acceptable salt thereof.

The invention also provides a pharmaceutical composition comprising a compound of the present invention and a pharmaceutically acceptable carrier.

The invention also provides a method of treating a tumor in a mammal in need of such treatment comprising administering to the mammal an

5 effective amount of a compound of the present invention; and administering  
neutron capture therapy.

10 The invention also provides a method for imaging a tumor in a  
mammal in need of such imaging comprising administering to the mammal a  
5 detectable amount of a compound of the present invention in combination with a  
pharmaceutically acceptable vehicle effective to image the tumor; and detecting  
the presence of the compound.

15 The invention also provides a compound of the present invention  
for use in medical therapy or diagnosis.

20 The invention also provides the use of a compound of the present  
invention for the manufacture of a medicament for imaging a tumor in a  
mammal (e.g., a human).

25 The invention also provides the use of a compound of the present  
invention for the manufacture of a medicament for treating a tumor in a mammal  
(e.g., a human).

30 The invention also provides intermediates disclosed herein that  
are useful in the preparation of the compounds of the present invention as well as  
synthetic methods useful for preparing the compounds of the invention.

#### **Brief Description of the Figures**

35 Figure 1 depicts the structure of cobalamin wherein X is CN  
(cyano), OH, CH<sub>3</sub> or adenosyl.

40 Figure 2 illustrates the synthesis of a cyanocobalamin-nido-  
carborane conjugate.

45 Figure 3 illustrates the synthesis of representative compounds of  
the present invention.

#### **Detailed Description of the Invention**

50 Applicant has discovered certain cobalamin analogs are useful, in  
combination with neutron capture therapy, to treat tumors. Applicant has also  
discovered certain cobalamin analogs are useful as neutron capture agents and as  
tumor imaging agents.

55 The following definitions are used, unless otherwise described:  
halo is fluoro, chloro, bromo, or iodo. Alkyl, alkoxy, alkenyl, alkynyl, etc.  
denote both straight and branched groups; but reference to an individual radical

5 such as "propyl" embraces only the straight chain radical, a branched chain isomer such as "isopropyl" being specifically referred to. Aryl denotes a phenyl radical or an ortho-fused bicyclic carbocyclic radical having about nine to ten ring atoms in which at least one ring is aromatic.

10 5 Specific and preferred values listed below for radicals, substituents, and ranges, are for illustration only; they do not exclude other defined values or other values within defined ranges for the radicals and substituents.

15 It is appreciated that those skilled in the art will recognize that  
10 compounds of the present invention having a chiral center may exist in and be isolated in optically active and racemic forms. Some compounds may exhibit polymorphism. It is to be understood that the present invention encompasses  
20 any racemic, optically-active, polymorphic, or stereoisomeric form, or mixtures thereof, of a compound of the invention, which possess the useful properties described herein, it being well known in the art how to prepare optically active  
25 forms (for example, by resolution of the racemic form by recrystallization techniques, by synthesis from optically-active starting materials, by chiral synthesis, or by chromatographic separation using a chiral stationary phase) and how to determine antitumor activity or utility as an antitumor imaging agent  
30 using the standard tests described herein, or using other similar tests which are well known in the art.

35 Specifically, (C<sub>1</sub>-C<sub>14</sub>)alkyl can be methyl, ethyl, propyl, isopropyl, butyl, iso-butyl, sec-butyl, pentyl, 3-pentyl, hexyl, heptyl, octyl, nonyl, decyl undecyl, dodecyl, tridecyl or tetradecyl.

25 Specifically, (C<sub>2</sub>-C<sub>14</sub>)alkenyl can be vinyl, allyl, 1-propenyl, 2-propenyl, 1-butenyl, 2-butenyl, 3-butenyl, 1-pentenyl, 2-pentenyl, 3-pentenyl, 4-pentenyl, 1-hexenyl, 2-hexenyl, 3-hexenyl, 4-hexenyl, 5-hexenyl, 1-heptenyl, 2-heptenyl, 3-heptenyl, 4-heptenyl, 5-heptenyl, 6-heptenyl, 1-octenyl, 2-octenyl, 3-octenyl, 4-octenyl, 5-octenyl, 6-octenyl, 7-octenyl, 1-nonenyl, 2-nonenyl, 3-  
40 nonenyl, 4-nonenyl, 5-nonenyl, 6-nonenyl, 7-nonenyl, 8-nonenyl, 1-decenyl, 2-decenyl, 3-decenyl, 4-decenyl, 5-decenyl, 6-decenyl, 7-decenyl, 8-decenyl, 9-decenyl, 1-undecenyl, 2-undecenyl, 3-undecenyl, 4-undecenyl, 5-undecenyl, 6-undecenyl, 7-undecenyl, 8-undecenyl, 9-undecenyl, 10-undecenyl, 1-dodecenyl,  
45  
50



2 -dodecenyl, 3 -dodecenyl, 4 -dodecenyl, 5-dodecenyl, 6-dodecenyl, 7-dodecenyl, 8-dodecenyl, 9-dodecenyl, 10-dodecenyl, 11-dodecenyl, 1-tridecenyl, 2-tridecenyl, 3-tridecenyl, 4-tridecenyl, 5-tridecenyl, 6-tridecenyl, 7-tridecenyl, 8-tridecenyl, 9-tridecenyl, 10-tridecenyl, 11-tridecenyl, 12-tridecenyl, 1-tetradecenyl, 2-tetradecenyl, 3-tetradecenyl, 4-tetradecenyl, 5-tetradecenyl, 6-tetradecenyl, 7-tetradecenyl, 8-tetradecenyl, 9-tetradecenyl, 10-tetradecenyl, 11-tetradecenyl, 12-tetradecenyl or 13-tetradecenyl.

Specifically, (C<sub>2</sub>-C<sub>14</sub>)alkynyl can be ethynyl, 1-propynyl, 2-propynyl, 1-butylnyl, 2-butylnyl, 3-butylnyl, 1-pentylnyl, 2-pentylnyl, 3-pentylnyl, 4-pentylnyl, 1-hexynyl, 2-hexynyl, 3-hexynyl, 4-hexynyl, 5-hexynyl, 1-heptylnyl, 2-heptylnyl, 3-heptylnyl, 4-heptylnyl, 5-heptylnyl, 6-heptylnyl, 1-octynyl, 2-octynyl, 3-octynyl, 4-octynyl, 5-octynyl, 6-octynyl, 7-octynyl, 1-nonylnyl, 2-nonylnyl, 3-nonylnyl, 4-nonylnyl, 5-nonylnyl, 6-nonylnyl, 7-nonylnyl, 8-nonylnyl, 1-decynyl, 2-decynyl, 3-decynyl, 4-decynyl, 5-decynyl, 6-decynyl, 7-decynyl, 8-decynyl, 9-decynyl, 1-undecynyl, 2-undecynyl, 3-undecynyl, 4-undecynyl, 5-undecynyl, 6-undecynyl, 7-undecynyl, 8-undecynyl, 9-undecynyl, 10-undecynyl, 1-dodecynyl, 2-dodecynyl, 3-dodecynyl, 4-dodecynyl, 5-dodecynyl, 6-dodecynyl, 7-dodecynyl, 8-dodecynyl, 9-dodecynyl, 10-dodecynyl, 11-dodecynyl, 1-tridecynyl, 2-tridecynyl, 3-tridecynyl, 4-tridecynyl, 5-tridecynyl, 6-tridecynyl, 7-tridecynyl, 8-tridecynyl, 9-tridecynyl, 10-tridecynyl, 11-tridecynyl, 12-tridecynyl, 1-tetradecynyl, 2-tetradecynyl, 3-tetradecynyl, 4-tetradecynyl, 5-tetradecynyl, 6-tetradecynyl, 7-tetradecynyl, 8-tetradecynyl, 9-tetradecynyl, 10-tetradecynyl, 11-tetradecynyl, 12-tetradecynyl or 13-tetradecynyl.

Specifically "aryl" can be phenyl, indenyl, or naphthyl.

Specifically (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl can be cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl or cyclooctyl.

As used herein, an "amino acid" is a natural amino acid residue (e.g. Ala, Arg, Asn, Asp, Cys, Glu, Gln, Gly, His, Hyl, Hyp, Ile, Leu, Lys, Met, Phe, Pro, Ser, Thr, Trp, Tyr, and Val) in D or L form, as well as unnatural amino acid (e.g. phosphoserine; phosphothreonine; phosphotyrosine; hydroxyproline; gamma-carboxyglutamate; hippuric acid; octahydroindole-2-carboxylic acid; statine; 1,2,3,4,-tetrahydroisoquinoline-3-carboxylic acid; penicillamine; ornithine; citruline; α-methyl-alanine; para-benzoylphenylalanine;

phenylglycine; propargylglycine; sarcosine; and tert-butylglycine) residue having one or more open valences. The term also comprises natural and unnatural amino acids bearing conventional amino protecting groups (e.g. acetyl, acyl, trifluoroacetyl, or benzyloxycarbonyl), as well as natural and unnatural amino acids protected at carboxy (e.g. as a (C<sub>1</sub>-C<sub>6</sub>)alkyl, phenyl or benzyl ester or amide) with conventional protecting groups. Other suitable amino and carboxy protecting groups are known to those skilled in the art (See for example, T.W. Greene, Protecting Groups In Organic Synthesis; Wiley: New York, 1981; D. Voet, Biochemistry, Wiley: New York, 1990; L. Stryer, Biochemistry, (3rd Ed.), W.H. Freeman and Co.: New York, 1975; J. March, Advanced Organic Chemistry, Reactions, Mechanisms and Structure, (2nd Ed.), McGraw Hill: New York, 1977; F. Carey and R. Sundberg, Advanced Organic Chemistry, Part B: Reactions and Synthesis, (2nd Ed.), Plenum: New York, 1977; and references cited therein). According to the invention, the amino or carboxy protecting group can also comprise a radionuclide (e.g., Fluorine-18, Iodine-123, or Iodine-124).

As used herein, a "peptide" is a sequence of 2 to 25 amino acids (e.g. as defined hereinabove) or peptidic residues having one or more open valences. The sequence may be linear or cyclic. For example, a cyclic peptide can be prepared or may result from the formation of disulfide bridges between two cysteine residues in a sequence. A peptide can be linked through the carboxy terminus, the amino terminus, or through any other convenient point of attachment, such as, for example, through the sulfur of a cysteine. Peptide derivatives can be prepared as disclosed in U.S. Patent Numbers 4,612,302; 4,853,371; and 4,684,620, or as described in the Examples hereinbelow. Peptide sequences specifically recited herein are written with the amino terminus on the left and the carboxy terminus on the right.

Particular and specific values listed below for radicals, substituents, and ranges, are for illustration only and they do not exclude other defined values or other values within defined ranges for the radicals and substituents.

Specifically, the peptide can be poly-L-lysine, poly-L-glutamic acid, poly-L-aspartic acid, poly-L-histidine, poly-L-ornithine, poly-L-serine,

poly-L-threonine, poly-L-tyrosine, poly-L-lysine--L-phenylalanine or poly-L-lysine-L-tyrosine.

As used herein, a "residue of a compound of formula I" is a radical of a compound of formula I having one or more open valences. Any synthetically feasible atom or atoms of the compound of formula I may be removed to provide the open valence, provided the resulting compound is able to localize in or near tumors. Based on the linkage that is desired, one skilled in the art can select suitably functionalized starting materials that can be derived from a compound of formula I using procedures that are known in the art. For example, suitable atoms that may be removed include the NH<sub>2</sub> group of the a-carboxamide (illustrated in figure 1) or a hydrogen atom from the NH<sub>2</sub> group of the a-carboxamide, the NH<sub>2</sub> group of the b-carboxamide (illustrated in figure 1) or a hydrogen atom from the NH<sub>2</sub> group of the b-carboxamide, the NH<sub>2</sub> group of the d-carboxamide (illustrated in figure 1) or a hydrogen atom from the NH<sub>2</sub> group of the d-carboxamide, the NH<sub>2</sub> group of the e-carboxamide (illustrated in figure 1) or a hydrogen atom from the NH<sub>2</sub> group of the e-carboxamide, and X at the 6-position (illustrated in figure 1). In addition, the hydrogen atom of the hydroxy group at the 3' position of the sugar, the hydrogen atom from the hydroxyl group at the 3' position of the sugar, the hydrogen atom of the CH<sub>2</sub>OH group at the 5' position, or the hydrogen atom from the hydroxyl group at the 5' position of the sugar ring may be removed.

As used herein, "adenosyl" is an adenosine radical in which any synthetically feasible atom or group of atoms have been removed, thereby providing an open valence. Synthetically feasible atoms which may be removed include the hydrogen atom of the hydroxy group at the 5' position. Accordingly, adenosyl can conveniently be attached to the 6-position of a compound of formula I via the 5' position of adenosyl.

As used herein, a "molecule comprising B-10" can be any compound that contains at least one B-10 atom. The nature of the molecule that includes B-10 is not critical. The compound, however, must be nontoxic and must be able to enter the tumor cell or locate near the tumor cell when the molecule comprising B-10 is attached.

As used herein, a "residue of a molecule comprising B-10" is a radical of a molecule comprising B-10 having one or more open valences. Any synthetically feasible atom or atoms of the molecule comprising B-10 may be removed to provide the open valence, provided bioactivity is substantially retained. Based on the linkage that is desired, one skilled in the art can select suitably functionalized starting materials that can be derived from a molecule comprising B-10 using procedures that are known in the art.

As used herein, a "residue of o-carborane," a "residue of m-carborane," or a "residue of p-carborane" is a radical of o-carborane, m-carborane or p-carborane, respectively, having one or more open valences. Any synthetically feasible atom or atoms of o-carborane, m-carborane or p-carborane may be removed to provide the open valence, provided bioactivity is substantially retained. Based on the linkage that is desired, one skilled in the art can select suitably functionalized starting materials that can be derived from o-carborane, m-carborane or p-carborane using procedures that are known in the art.

#### Boron Compounds Useful For Neutron Capture Therapy

The invention provides a compound of formula I (Figure 1) linked to one or more molecules comprising B-10, wherein X is CN, OH, CH<sub>3</sub>, adenosyl or a molecule comprising B-10; or a pharmaceutically acceptable salt thereof.

A variety of molecules comprising B-10 known in the art are useful in the present invention. The molecules vary considerably in structure but are suitable to practice the present invention. Acceptable species include boron containing amino acids, carbohydrates, and nucleosides, as well as carboranes. A wide variety of boron-containing compounds are commercially available, or are known in the art. A variety of molecules comprising B-10 are commercially available from Boron Biologicals, Inc., Raleigh, North Carolina and RysCor Science, Inc., Raleigh, North Carolina.

Specifically, at least one molecule comprising B-10 can be o-carborane, m-carborane or p-carborane. More specifically, at least one molecule comprising B-10 is o-carborane. o-Carborane [1,2-dicarbadoecaborane(12)]; m-carborane [1,7-dicarbadoecaborane(12)]; and p-carborane [1,12-

dicarbadoecaborane(12)] are commercially available from Aldrich, Milwaukee, WI.

Specifically, each molecule comprising B-10 can independently be o-carborane, m-carborane or p-carborane. More specifically, each molecule comprising B-10 is o-carborane.

Compound of Formula I / Molecule Comprising B-10

The residue of molecule comprising B-10 can be linked to the residue of a compound of formula I through an amide (e.g., -N(R)C(=O)- or -C(=O)N(R)-), ester (e.g., -OC(=O)- or -C(=O)O-), ether (e.g., -O-), amino (e.g., -N(R)-), ketone (e.g., -C(=O)-), thioether (e.g., -S-), sulfinyl (e.g., -S(O)-), sulfonyl (e.g., -S(O)<sub>2</sub>-), or a direct (e.g., C-C bond) linkage, wherein each R is independently H or (C<sub>1</sub>-C<sub>6</sub>)alkyl. Such a linkage can be formed from suitably functionalised starting materials using synthetic procedures that are known in the art. Based on the linkage that is desired, one skilled in the art can select suitably functional starting materials that can be derived from a residue of a compound of formula I and from a given residue of a molecule comprising B-10 using procedures that are known in the art.

The residue of the molecule comprising B-10 can be directly linked to any synthetically feasible position on the residue of a compound of formula I. Suitable points of attachment include, for example, the b-carboxamide, the d-carboxamide, and the e-carboxamide (illustrated in figure 1), as well as the 6-position (the position occupied by X in figure 1), and the 5'-hydroxy and the 3'-hydroxy groups on the 5-membered sugar ring, although other points of attachment are possible. U.S. Patent No. 5,739,313 discloses compounds (e.g., cyanocobalamin-b-(4-aminobutyl)amide, methylcobalamin-b-(4-aminobutyl)amide, and adenosylcobalamin-b-(4-aminobutyl)amide) that are useful intermediates for the preparation of compounds of the present invention.

Compounds wherein the residue of a molecule comprising B-10 is linked to the 6-position of a compound of formula I can be prepared by reducing a corresponding Co (II) compound of formula I to form a nucleophilic Co (I) compound and treating this Co (I) compound with a residue of a molecule comprising B-10 (or a derivative thereof) comprising a suitable leaving group, such as a halide (e.g., a chloride).

5 The invention also provides compounds having more than one  
residue of a molecule comprising B-10 directly linked to a compound of formula  
I. For example, the residue of a molecule comprising B-10 can be directly linked  
10 to a residue of the b-carboxamide of the compound of formula I and a residue of  
5 another molecule comprising B-10 can be directly linked to a residue of the d-  
carboxamide of the compound of formula I. In addition, the residue of a  
molecule comprising B-10 can be directly linked to the 6-position of the  
15 compound of formula I and a residue of another molecule comprising B-10 can  
be directly linked to a residue of the d- or e-carboxamide of the compound of  
10 formula I.

Compound of Formula I / Linker / Molecule Comprising B-10 Linker

20 In addition to being directly linked to the residue of a compound  
of formula I, the residue of a molecule comprising B-10 can also be linked to the  
residue of a compound of formula I by a suitable linker. The structure of the  
15 linker is not crucial, provided it yields a compound of the invention which has an  
effective therapeutic index against the target cells, and which will localize in or  
25 near tumor molecules.

Suitable linkers include linkers that separate the residue of a  
30 compound of formula I and the residue of a molecule comprising B-10 by about  
20 5 angstroms to about 200 angstroms. Other suitable linkers include linkers that  
separate the residue of a compound of formula I and the residue of a molecule  
comprising B-10 by about 5 angstroms to about 100 angstroms, as well as  
35 linkers that separate the residue of a compound of formula I and the residue of a  
molecule comprising B-10 by about 5 angstroms to about 50 angstroms, or by  
25 about 5 angstroms to about 25 angstroms. Suitable linkers are disclosed, for  
example, in U.S. Patent No. 5,735,313.

40 The linker can be linked to any synthetically feasible position on  
the residue of a compound of the residue of formula I. Suitable points of  
attachment include, for example, a residue of the b-carboxamide, a residue of the  
45 d-carboxamide, and a residue of the e-carboxamide, the 6-position (i.e., the  
30 position occupied by X in the compound of formula I), as well as a residue of the  
5'-hydroxy group and a residue of the 3'-hydroxy group on the 5-membered  
50 sugar ring, although other points of attachment are possible. Based on the

linkage that is desired, one skilled in the art can select suitably functionalized starting materials that can be derived from a compound of formula I and from a given molecule comprising B-10 using procedures that are known in the art.

The linker can conveniently be linked to the residue of a compound of formula I or to the residue of a molecule comprising B-10 through an amide (e.g., -N(R)C(=O)- or -C(=O)N(R)-), ester (e.g., -OC(=O)- or -C(=O)O-), ether (e.g., -O-), ketone (e.g., -C(=O)-) thioether (e.g., -S-), sulfinyl (e.g., -S(O)-), sulfonyl (e.g., -S(O)<sub>2</sub>-), amino (e.g., -N(R)-) or a direct (e.g., C-C) linkage, wherein each R is independently H or (C<sub>1</sub>-C<sub>6</sub>)alkyl. The linkage can be formed from suitably functionalised starting materials using synthetic procedures that are known in the art. Based on the linkage that is desired, one skilled in the art can select suitably functional starting materials that can be derived from a residue of a compound of formula I, a residue of a molecule comprising B-10, and from a given linker using procedures that are known in the art.

Specifically, the linker can be a divalent radical of the formula W-A-Q wherein A is (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>2</sub>-C<sub>6</sub>)alkynyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, or (C<sub>6</sub>-C<sub>10</sub>)aryl, wherein W and Q are each independently -N(R)C(=O)-, -C(=O)N(R)-, -OC(=O)-, -C(=O)O-, -O-, -S-, -S(O)-, -S(O)<sub>2</sub>-, -N(R)-, -C(=O)-, or a direct bond (i.e., W or Q is absent); wherein each R is independently H or (C<sub>1</sub>-C<sub>6</sub>)alkyl.

Specifically, the linker can be a divalent radical of the formula W-(CH<sub>2</sub>)<sub>n</sub>-Q wherein, n is between about 1 and about 20, between about 1 and about 15, between about 2 and about 10, between about 2 and about 6, or between about 4 and about 6; wherein W and Q are each independently -N(R)C(=O)-, -C(=O)N(R)-, -OC(=O)-, -C(=O)O-, -O-, -S-, -S(O)-, -S(O)<sub>2</sub>-, -C(=O)-, -N(R)-, or a direct bond (i.e., W or Q is absent); wherein each R is independently H or (C<sub>1</sub>-C<sub>6</sub>)alkyl.

Specifically, W and Q are each independently -N(R)C(=O)-, -C(=O)N(R)-, -OC(=O)-, -N(R)-, -C(=O)O-, -O-, or a direct bond (i.e., W or Q is absent).

Specifically, the linker is a divalent radical, i.e., 1,ω-divalent radicals formed from a peptide or an amino acid. The peptide can comprise 2 to about 20 amino acids, 2 to about 15 amino acids, or 2 to about 12 amino acids.

Specifically, the peptide can be poly-L-lysine (i.e.,  $[-NHCH[(CH_2)_4NH_2]CO-]_m-Q$ , wherein Q is H,  $(C_1-C_{14})$ alkyl, or a suitable carboxy protecting group; and wherein m is about 2 to about 20. Specifically, the poly-L-lysine can contain about 5 to about 15 residues (i.e., m is between about 5 and about 15). More specifically, the poly-L-lysine can contain about 8 to about 11 residues (i.e., m is between about 8 and about 11).

Specifically, the peptide can be poly-L-glutamic acid, poly-L-aspartic acid, poly-L-histidine, poly-L-ornithine, poly-L-serine, poly-L-threonine, poly-L-tyrosine, poly-L-lysine-L-phenylalanine or poly-L-lysine-L-tyrosine.

Specifically, the linker is prepared from 1,6-diaminohexane  $H_2N(CH_2)_6NH_2$ , 1,5-diaminopentane  $H_2N(CH_2)_5NH_2$ , 1,4-diaminobutane  $H_2N(CH_2)_4NH_2$ , or 1,3-diaminopropane  $H_2N(CH_2)_3NH_2$ .

The linker can comprise one or more non-metallic radionuclides. Specifically, the linker can comprise more than one non-metallic radionuclides. More specifically, the linker can comprise 2 to about 10, 2 to about 8, 2 to about 6, or 2 to about 4 non-metallic radioisotopes.

A specific residue of a peptide (i.e., linker) comprising one or more non-metallic radionuclides has the following formula


wherein each M is

independently a non-metallic radionuclide; each R is independently  $(C_1-C_{14})$ alkyl,  $(C_2-$

$C_{14})$ alkenyl,  $(C_2-C_{14})$ alkynyl,  $(C_1-$

$C_{14})$ alkoxy, hydroxy, cyano, nitro, halo, trifluoromethyl,  $N(R_a)(R_b)$ ,  $(C_1-C_{14})$ alkanoyl,  $(C_2-C_{14})$ alkanoyloxy,  $(C_6-C_{10})$ aryl, or  $(C_3-C_8)$ cycloalkyl wherein  $R_a$  and  $R_b$  are each independently H or  $(C_1-C_{14})$ alkyl; P; Q is H,  $(C_1-C_{14})$ alkyl, or a suitable carboxy protecting group; n is 2 to about 20; I is 1-5, j is 0-4 and  $I+j \leq 5$ ; or a pharmaceutically acceptable salt thereof.

Specifically, i can be 1, j can be 0, M can be Fluorine-18, Bromine-76, or Iodine-123, and n can be about 6 to about 12.





5 The molecule comprising B-10 can comprise one or more boron atoms. A suitable molecule comprising B-10 can contain 1 to about 20, 1 to about 15, 1 to about 10, or 1 to about 5 boron atoms. In addition, the molecule comprising B-10 can comprise one or more B-10 atoms. A suitable molecule  
10 5 comprising B-10 can contain 1 to about 20, 1 to about 15, 1 to about 10, or 1 to about 5 B-10 atoms.

The molecule comprising B-10 can be an amino acid, a carbohydrate, a nucleoside or a carborane. Specifically, the molecule comprising  
15 B-10 is o-carborane, m-carborane or p-carborane.

10 Compounds wherein the linker is linked to the 6-position of a compound of formula I can be prepared by preparing a nucleophilic Co (I) species as described herein above, and reacting it with a linker comprising a  
20 suitable leaving group, such as a halide (e.g. a chloride).

The invention also provides compounds having more than one  
25 molecule comprising B-10 attached to a compound of formula I, each through a linker. For example, the residue of a molecule comprising B-10 can conveniently be linked, through a linker, to a residue of the b-carboxamide of the compound of formula I and a residue of another molecule comprising B-10 can  
30 conveniently be linked, through a linker, to a residue of the d- or e-carboxamide of the compound of formula I. In addition, the residue of a molecule comprising B-10 can conveniently be linked, through a linker, to the 6-position of the compound of formula I and a residue of another molecule comprising B-10 can  
35 conveniently be linked, through a linker, to a residue of the b-, d- or e-carboxamide of the compound of formula I.

25 The invention also provides compounds having more than one molecule comprising B-10 attached to a compound of formula I, either directly  
40 or through a linker. For example, the residue of a molecule comprising B-10 can conveniently be linked, either directly or through a linker, to a residue of the b-carboxamide of the compound of formula I and a residue of another molecule  
45 comprising B-10 can conveniently be linked, either directly or through a linker, to a residue of the d- or e-carboxamide of the compound of formula I. In addition, the residue of a molecule comprising B-10 can conveniently be linked,  
50 either directly or through a linker, to the 6-position of the compound of formula I

5 and a residue of another molecule comprising B-10 can conveniently be linked, either directly or through a linker, to a residue of the b-, d- or e-carboxamide of the compound of formula I.

Compound of Formula I / Chelating Group Comprising Gadolinium-157

10 5 U.S. Patent Number 5,739,313 discloses cobalamin analogs comprising a compound of formula I, a linking group, and a chelating group comprising a detectable radionuclide or a paramagnetic ion. The compounds are disclosed to localize in tumor cells following administration, and to be useful for  
15 imaging tumors.

10 The metallic radionuclide Gadolinium-157 is an especially useful ion for conducting magnetic resonance imaging. It is also a useful target ion for  
20 neutron capture therapy. Applicants have discovered that incorporation of Gd-157 into one of the chelating molecules disclosed in U.S. Patent Number 5,739,313, provides a compound that is not only particularly useful for  
25 conducting magnetic resonance imaging, but also a compound that can be used in conjunction with neutron capture therapy, to treat tumors.

Thus, the present invention provides a residue of a compound of formula I linked to at least one residue of the formula -Q-L-W-Det; wherein each  
30 Det is independently a chelating group comprising Gd-157; each L is independently linker (as defined hereinabove) or is absent; and each W and Q are  
20 each independently -N(R)C(=O)-, -C(=O)N(R)-, -OC(=O)-, -C(=O)O-, -O-, -S-, -S(O)-, -S(O)<sub>2</sub>-, -C(=O)-, -N(R)-, or a direct bond (i.e., W or Q is absent);  
35 wherein each R is independently H or (C<sub>1</sub>-C<sub>6</sub>)alkyl.

Compound of Formula I / Chelating Group Comprising a Radionuclide

25 Applicant has also discovered that it is possible to prepare a compound that is useful for both imaging and for treating tumors by  
40 incorporating one or more neutron capture target atoms (e.g. B-10 or Gd-157) into a compound that also comprises a detectable radionuclide. Accordingly, the invention provides a residue of a compound of formula I which is linked to one  
45 30 or more residues of a molecule comprising B-10; and which is also linked to one or more groups of the formula -Q-L-W-Det; wherein each Det is independently a chelating group (as defined herein) comprising Gd-157; each L is independently linker (as defined herein) or is absent; and each W and Q are each independently  
50

5 -N(R)C(=O)-, -C(=O)N(R)-, -OC(=O)-, -C(=O)O-, -O-, -S-, -S(O)-, -S(O)<sub>2</sub>-, -C(=O)-, -N(R)-, or a direct bond (i.e., W or Q is absent); wherein each R is independently H or (C<sub>1</sub>-C<sub>6</sub>)alkyl; or a pharmaceutically acceptable salt thereof.

10 The present invention provides a residue of a compound of formula I linked to a molecule comprising B-10 or linked to a chelating group comprising Gd-157 and wherein the residue of a compound of formula I is linked to at least one residue of the formula -Q-L-W-Det; wherein each Det is independently a chelating group (as defined hereinabove) comprising a metallic radionuclide; each L is independently a linker (as defined hereinabove) or is  
15 absent; and each W and Q are each independently -N(R)C(=O)-, -C(=O)N(R)-, -OC(=O)-, -C(=O)O-, -O-, -S-, -S(O)-, -S(O)<sub>2</sub>-, -C(=O)-, -N(R)-, or a direct bond (i.e., W or Q is absent); wherein each R is independently H or (C<sub>1</sub>-C<sub>6</sub>)alkyl.  
20

Compound of Formula I / Molecule Comprising B-10 / Detectable Radionuclide.

15 The invention also provides a compound wherein a residue of a compound of formula I (Figure 1) is linked 1) to a residue of a molecule comprising B-10, and 2) to a detectable radionuclide, wherein X is CN, OH, CH<sub>3</sub>, adenosyl, or a molecule comprising B-10; or a pharmaceutically acceptable salt thereof.  
25

20 The detectable radionuclide can be directly linked to the residue of a compound of formula I, or can be linked by a linker to the residue of a compound of formula I. The linker can be any suitable linker described herein. Specifically, the linker can be of the formula W-A wherein A is (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>2</sub>-C<sub>6</sub>)alkynyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, or (C<sub>6</sub>-C<sub>10</sub>)aryl, wherein W is  
35 -N(R)C(=O)-, -C(=O)N(R)-, -OC(=O)-, -C(=O)O-, -O-, -S-, -S(O)-, -S(O)<sub>2</sub>-, -N(R)-, -C(=O)-, or a direct bond; wherein each R is independently H or (C<sub>1</sub>-C<sub>6</sub>)alkyl; and wherein A is substituted with one or more non-metallic radionuclides. The linker can also be a peptide or an amino acid.  
40

Compound of Formula I / Group comprising Gd-157 / Detectable Radionuclide.

The invention also provides a compound wherein a residue of a compound of formula I (Figure 1) is linked 1) to a detectable radionuclide; and 2) to a group comprising Gd-157; or a pharmaceutically acceptable salt thereof.

The Gd-157 can be linked to the residue of a compound of formula I by any suitable means. For example, the residue of a compound of formula I can be attached to a group comprising Gd-157 that has the formula Q-L-W-Det, wherein Det is a chelating group comprising Gd-157, L is a linker or absent, and W and Q are each independently -N(R)C(=O)-, -C(=O)N(R)-, -OC(=O)-, -C(=O)O-, -O-, -S-, -S(O)-, -S(O)<sub>2</sub>-, -C(=O)-, -N(R)-, or a direct bond; wherein each R is independently H or (C<sub>1</sub>-C<sub>6</sub>)alkyl. Any suitable chelator can be used.

The detectable radionuclide can be directly linked to the residue of a compound of formula I, or can be linked by a linker to the residue of a compound of formula I. The linker can be any suitable linker described herein. Specifically, the linker can be of the formula W-A wherein A is (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>2</sub>-C<sub>6</sub>)alkynyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, or (C<sub>6</sub>-C<sub>10</sub>)aryl, wherein W is -N(R)C(=O)-, -C(=O)N(R)-, -OC(=O)-, -C(=O)O-, -O-, -S-, -S(O)-, -S(O)<sub>2</sub>-, -N(R)-, -C(=O)-, or a direct bond; wherein each R is independently H or (C<sub>1</sub>-C<sub>6</sub>)alkyl; and wherein A is substituted with one or more non-metallic radionuclides. The linker can also be a peptide or an amino acid

Non-Metallic Radionuclide

Any detectable non-metallic radionuclide that is suitable for imaging can be used in the compounds of the invention. For example, suitable non-metallic radionuclides include Carbon-11, Fluorine-18, Bromine-76, and Iodine-123. Specifically, the non-metallic radionuclide can be a non-metallic paramagnetic atom (e.g., Fluorine-19); or a non-metallic positron emitting radionuclide (e.g., Carbon-11, Fluorine-18, Iodine-123, or Bromine-76).

Metallic Radionuclide

Suitable metallic radionuclides (i.e., metallic radioisotopes or metallic paramagnetic ions) include Antimony-124, Antimony-125, Arsenic-74, Barium-103, Barium-140, Beryllium-7, Bismuth-206, Bismuth-207, Cadmium-109, Cadmium-115m, Calcium-45, Cerium-139, Cerium-141, Cerium-144,

5 Cesium-137, Chromium-51, Cobalt-55, Cobalt-56, Cobalt-57, Cobalt-58,  
Cobalt-60, Cobalt-64, Copper-67, Erbium-169, Europium-152, Gallium-64,  
Gallium-68, Gadolinium-153, Gadolinium-157 Gold-195, Gold-199, Hafnium-  
10 175, Hafnium-175-181, Holmium-166, Indium-110, Indium-111, Iridium-192,  
5 Iron-55, Iron-59, Krypton-85, Lead-210, Manganese-54, Mercury-197, Mercury-  
203, Molybdenum-99, Neodymium-147, Neptunium-237, Nickel-63, Niobium-  
95, Osmium-185 + 191, Palladium-103, Platinum-195m, Praseodymium-143,  
15 Promethium-147, Protactinium-233, Radium-226, Rhenium-186, Rhenium-188,  
Rubidium-86, Ruthenium-103, Ruthenium-106, Scandium-44, Scandium-46,  
10 Selenium-75, Silver-110m, Silver-111, Sodium-22, Strontium-85, Strontium-89,  
Strontium-90, Sulfur-35, Tantalum-182, Technetium-99m, Tellurium-125,  
20 Tellurium-132, Thallium-204, Thorium-228, Thorium-232, Thallium-170, Tin-  
113, Tin-114, Tin-117m, Titanium-44, Tungsten-185, Vanadium-48, Vanadium-  
49, Ytterbium-169, Yttrium-86, Yttrium-88, Yttrium-90, Yttrium-91, Zinc-65,  
15 and Zirconium-95.


25 As used herein, a "detectable radionuclide" is any suitable  
radionuclide (i.e., radioisotope) capable of detecting cancer or other neoplastic  
cells in a diagnostic procedure *in vivo* or *in vitro*. Suitable detectable  
30 radionuclides include metallic radionuclides (i.e., metallic radioisotopes) and  
20 non-metallic radionuclides (i.e., non-metallic radioisotopes).

As used herein, a "therapeutic radionuclide" is any suitable  
radionuclide (i.e., radioisotope) that possesses therapeutic efficacy against cancer  
35 or other neoplastic cells *in vivo* or *in vitro*. Suitable therapeutic radionuclides  
include metallic radionuclides (i.e., metallic radioisotopes).

25 Chelating groups

40 Any suitable chelating group can be incorporated into the  
compounds of the invention. Suitable chelating groups include those disclosed  
in U.S. Patent Number 5,739,313. Specifically, the chelating group can be NTA,  
HEDTA, DCTA, RP414, MDP, DOTATOC, CDTA, HYNIC, EDTA, DTPA,  
45 30 TETA, DOTA, DOTMP, DCTA, 15N4, 9N3, 12N3, or MAG3 (or another  
suitable polyamino acid chelator), which are described herein below, or a  
phosphonate chelator (e.g. EDTM). More specifically, the chelating group can  
50 be DTPA.

DTPA is diethylenetriaminepentaacetic acid; TETA is 1,4,8,11-tetraazacyclotetradecane-N,N',N'',N'''-tetraacetic acid; DOTA is 1,4,7,10-tetraazacyclododecane-N,N',N'',N'''-tetraacetic acid; 15N4 is 1,4,8,12-tetraazacyclopentadecane-N,N',N'',N'''-tetraacetic acid; 9N3 is 1,4,7-triazacyclononane-N,N',N''-triacetic acid; 12N3 is 1,5,9-triazacyclododecane-N,N',N''-triacetic acid; MAG3 is (N-[N-[N-[(benzoylthio)acetyl]glycyl]glycyl]glycine); and DCTA is a cyclohexane-based metal chelator of the formula



wherein R<sup>3</sup> may be (C<sub>1</sub>-C<sub>4</sub>)alkyl or CH<sub>2</sub>CO<sub>2</sub><sup>-</sup>, which may be attached through positions 4 or 5, or through the group R<sup>3</sup> and which carries from 1 to 4 detectable metal or nonmetal cations (M), monovalent cations, or the alkaline earth metals. Thus, with metals of oxidation state +1, each individual cyclohexane-based molecule may carry up to 4 metal cations (where both R<sup>3</sup> groups are CH<sub>2</sub>COOM). As is more likely, with higher oxidation states, the number of metals will decrease to 2 or even 1 per cyclohexane skeleton. This formula is not intended to limit the molecule to any specific stereochemistry.

NTA, HEDTA, and DCTA are disclosed in Poster Sessions, Proceedings of the 46th Annual Meeting, J. Nuc. Med., p. 316, No. 1386. RP414 is disclosed in Scientific Papers, Proceedings of the 46th Annual Meeting, J. Nuc. Med., p. 123, No. 499. MDP is disclosed in Scientific Papers, Proceedings of the 46th Annual Meeting, J. Nuc. Med., p. 102, No. 413. DOTATOC is disclosed in Scientific Papers, Proceedings of the 46th Annual Meeting, J. Nuc. Med., p. 102, No. 414 and Scientific Papers, Proceedings of the 46th Annual Meeting, J. Nuc. Med., p. 103, No. 415. CDTA is disclosed in Poster Sessions, Proceedings of the 46th Annual Meeting, J. Nuc. Med., p. 318, No.

1396. HYNIC is disclosed in Poster Sessions, Proceedings of the 46th Annual Meeting, J. Nuc. Med., p. 319, No. 1398.

Bifunctional chelators (i.e., chelating groups) based on macrocyclic ligands in which conjugation is via an activated arm attached to the carbon backbone of the ligand can also be employed as a chelating group, as described by M. Moi et al., J. Amer. Chem. Soc., 49, 2639 (1989) (2-p-nitrobenzyl-1,4,7,10-tetraazacyclododecane-N,N',N'',N'''-tetraacetic acid); S. V. Deshpande et al., J. Nucl. Med., 31, 473 (1990); G. Kuser et al., Bioconj. Chem., 1, 345 (1990); C. J. Broan et al., J. C. S. Chem. Comm., 23, 1739 (1990); and C. J. Anderson et al., J. Nucl. Med. 36, 850 (1995) (6-bromoacetamido-benzyl-1,4,8,11-tetraazacyclotetradecane-N,N',N'',N'''-tetraacetic acid (BAT)).

In addition, the diagnostic chelator or therapeutic chelating groups can be any of the chelating groups disclosed in Scientific Papers, Proceedings of the 46th Annual Meeting, J. Nuc. Med., Wednesday, June 9, 1999, p. 124, No. 500.

A "detectable chelating group" is a chelating group comprising a metallic radionuclide (e.g., a metallic radioisotope) capable of detecting cancer or other neoplastic cells *in vivo* or *in vitro*.

Specifically, the chelating group can be any one of the carbonyl complexes disclosed in Waibel et al., Nature Biotechnology, 897-901, Vol. 17, September 1999; or Sattelberger et al., Nature Biotechnology, 849-850, Vol. 17, September 1999.

Specifically, the detectable chelating group can be any of the carbonyl complexes disclosed in Waibel et al., Nature Biotechnology, 897-901, Vol. 17, September 1999; or Sattelberger et al., Nature Biotechnology, 849-850, Vol. 17, September 1999, further comprising a metallic radionuclide. More specifically, the detectable chelating group can be any of the carbonyl complexes disclosed in Waibel et al., Nature Biotechnology, 897-901, Vol. 17, September 1999; or Sattelberger et al., Nature Biotechnology, 849-850, Vol. 17, September 1999, further comprising Technetium-99m.

As used herein, a "therapeutic chelating group" is a chelating group comprising a metallic radionuclide (e.g., a metallic radioisotope) that

5 possesses therapeutic efficacy against cancer or other neoplastic cells *in vivo* or  
10 *in vitro*. Any suitable chelating group can be employed.

Specifically, the therapeutic chelating group can be any of the  
carbonyl complexes disclosed in Waibel et al., *Nature Biotechnology*, 897-901,  
10 5 Vol. 17, September 1999; or Sattelberger et al., *Nature Biotechnology*, 849-850,  
Vol. 17, September 1999, further comprising a metallic radionuclide. More  
specifically, the therapeutic chelating group can be any of the carbonyl  
15 complexes disclosed in Waibel et al., *Nature Biotechnology*, 897-901, Vol. 17,  
September 1999; or Sattelberger et al., *Nature Biotechnology*, 849-850, Vol. 17,  
20 September 1999, further comprising Rhenium-186 or Rhenium-188.

Tumors treatable with the compounds and methods of the  
20 invention can be located in any part of the mammal. Specifically, the tumor can  
be located in the breast, lung, thyroid, lymph node, genitourinary system (e.g.,  
kidney, ureter, bladder, ovary, teste, or prostate), musculoskeletal system (e.g.,  
25 bones, skeletal muscle, or bone marrow), gastrointestinal tract (e.g., stomach,  
esophagus, small bowel, colon, rectum, pancreas, liver, or smooth muscle),  
central or peripheral nervous system (e.g., brain, spinal cord, or nerves), head  
and neck tumors (e.g., ears, eyes, nasopharynx, oropharynx, or salivary glands),  
30 or the heart.

20 The compounds disclosed herein can be prepared using  
procedures similar to those described in U.S. Patent Number 5,739,313, or using  
procedures similar to those described herein. The residue of a molecule  
35 comprising B-10 can be linked to the residue of a compound of formula I as  
described hereinabove. Additional intermediates and synthetic procedures useful  
25 for preparing compounds of the invention are disclosed, for example, in  
Hogenkamp, H. et al., *Synthesis and Characterization of nido-Carborane-*  
40 *Cobalamin Conjugates*, Nucl. Med. & Biol., 2000, 27, 89-92; Collins, D., et al.,  
*Tumor Imaging Via Indium 111-Labeled DTPA-Adenosylcobalamin*, Mayo  
Clinic Proc., 1999, 74:687-691; U.S. Application Ser. No. 60/129,733 filed 16  
45 30 April 1999; U.S. Application Ser. No. 60/159,874 filed 15 October 1999; U.S.  
Application Ser. No. 60/159,753 filed 15 October 1999; U.S. Application Ser.  
No. 60/159,873 filed 15 October 1999; and references cited therein.



5 In cases where compounds are sufficiently basic or acidic to form  
stable nontoxic acid or base salts, administration of the compounds as salts may  
be appropriate. Examples of pharmaceutically acceptable salts are organic acid  
addition salts formed with acids which form a physiological acceptable anion,  
10 5 for example, tosylate, methanesulfonate, acetate, citrate, malonate, tartarate,  
succinate, benzoate, ascorbate,  $\alpha$ -ketoglutarate, and  $\alpha$ -glycerophosphate.  
Suitable inorganic salts may also be formed, including hydrochloride, sulfate,  
nitrate, bicarbonate, and carbonate salts.

15 Pharmaceutically acceptable salts may be obtained using standard  
10 procedures well known in the art, for example by reacting a sufficiently basic  
compound such as an amine with a suitable acid affording a physiologically  
acceptable anion. Alkali metal (for example, sodium, potassium or lithium) or  
20 alkaline earth metal (for example calcium) salts of carboxylic acids can also be  
made.

25 The compounds of the present invention can be formulated as  
pharmaceutical compositions and administered to a mammalian host, such as a  
human patient in a variety of forms adapted to the chosen route of  
administration, i.e., orally or parenterally (e.g., by intravenous, intramuscular,  
30 intraperitoneal). Preferably, the compounds are administered parenterally.

20 The active compound may also be administered intravenously or  
intraperitoneally by infusion or injection. Solutions of the active compound or  
its salts can be prepared in water, optionally mixed with a nontoxic surfactant.  
35 Dispersions can also be prepared in glycerol, liquid polyethylene glycols,  
triacetin, and mixtures thereof and in oils. Under ordinary conditions of storage  
25 and use, these preparations contain a preservative to prevent the growth of  
microorganisms.

40 The pharmaceutical dosage forms suitable for injection or  
infusion can include sterile aqueous solutions or dispersions or sterile powders  
comprising the active ingredient which are adapted for the extemporaneous  
45 30 preparation of sterile injectable or infusible solutions or dispersions, optionally  
encapsulated in liposomes. In all cases, the ultimate dosage form should be  
sterile, fluid and stable under the conditions of manufacture and storage. The  
liquid carrier or vehicle can be a solvent or liquid dispersion medium

5 comprising, for example, water, ethanol, a polyol (for example, glycerol,  
propylene glycol, liquid polyethylene glycols, and the like), vegetable oils,  
nontoxic glyceryl esters, and suitable mixtures thereof. The proper fluidity can  
be maintained, for example, by the formation of liposomes, by the maintenance  
10 5 of the required particle size in the case of dispersions or by the use of surfactants.  
The prevention of the action of microorganisms can be brought about by various  
antibacterial and antifungal agents, for example, parabens, chlorobutanol,  
phenol, sorbic acid, thimerosal, and the like. In many cases, it will be preferable  
15 to include isotonic agents, for example, sugars, buffers or sodium chloride.  
10 Prolonged absorption of the injectable compositions can be brought about by the  
use in the compositions of agents delaying absorption, for example, aluminum  
monostearate and gelatin.  
20

Sterile injectable solutions are prepared by incorporating the  
active compound in the required amount in the appropriate solvent with various  
15 15 of the other ingredients enumerated above, as required, followed by filter  
sterilization. In the case of sterile powders for the preparation of sterile  
injectable solutions, the preferred methods of preparation are vacuum drying and  
the freeze drying techniques, which yield a powder of the active ingredient plus  
30 any additional desired ingredient present in the previously sterile-filtered  
20 solutions.

For illustration, suitable doses of a compound of the invention for  
use in therapy, in conjunction with neutron capture, include doses in the range of  
35 from about 0.1  $\mu\text{g}$  to about 100  $\mu\text{g}$ , e.g., from about 0.5  $\mu\text{g}$  to about 50  $\mu\text{g}$ , or  
from about 0.5  $\mu\text{g}$  to 15  $\mu\text{g}$  per treatment. Suitable doses for use in imaging or  
25 for use in imaging and therapy include doses in the range of from about 0.1 mg  
to about 50 g, e.g., from about .5 mg to about 10 g, or from about 0.5 g to 2 g per  
40 treatment.

The desired dose may conveniently be presented in a single dose  
or as divided doses administered at appropriate intervals, for example, as two,  
45 30 three, four or more sub-doses per day. The sub-dose itself may be further  
divided, e.g., into a number of discrete loosely spaced administrations.

The compounds are preferably dissolved or dispersed in a  
nontoxic liquid vehicle, such as physiological saline or a similar aqueous  
50

5 vehicle, to the desired concentration. A preselected therapeutic unit dose is then  
administered to the test animal or human patient, by oral administration or  
ingestion or by parenteral administration, as by intravenous or intraperitoneal  
infusion or injection, to attain the desired *in vivo* concentration. Doses useful for  
10 5 treating tumors can be derived, from those found to be effective to treat tumors  
in humans *in vitro* or in animal models, such as those described hereinbelow, or  
from dosages of other labeled vitamin B<sub>12</sub> molecules, previously employed in  
animal therapy.

15 Figure 2 and Figure 3 each illustrate the four step synthesis used  
10 to prepare three cyanocobalamin-nido-carborane conjugates. o-Carborane  
carboxylic acid (2) was prepared by reacting o-carborane with n-butyllithium  
and carbondioxide in ether for approximately one hour at -78°C. Treatment with  
20 thionyl chloride gave o-carborane carboxylic acid chloride (3), which was  
allowed to react with 1,4-butanediamine in pyridine to give the amide linked  
15 nido-carboranoyl(4-aminobutyl)amide (4). Compound (4) was linked to the b-  
monocarboxylic acid of cyanocobalamin, the d-monocarboxylic acid of  
cyanocobalamin, or both the b- and d-dicarboxylic acid of cyanocobalamin. In  
each reaction, hydroxybenzotriazole and 1-ethyl-3-(3-dimethylaminopropyl)  
25 carbodiimide was added to facilitate the formation of the amide bond.

30 The invention may be further illustrated by the following  
examples.

### Examples

35 Ultraviolet-visible spectra were recorded with a diode array  
spectrophotometer. <sup>11</sup>B NMR spectra at 192.656 MHz with <sup>1</sup>H decoupling were  
25 recorded on Varian 1NOVA 600 MHz spectrometer with an 8 mm broad band  
probe. Approximately 10 mg of the cobalamin-conjugate were dissolved in 1  
40 mL pyridine d<sub>6</sub> in a 5 mm NMR tube. Five hundred scans were collected with an  
acquisition time of 0.133 seconds and a relaxation delay of 1 second. Chemical  
shifts are given relative to BF<sub>3</sub>·(CH<sub>3</sub>CH<sub>2</sub>)<sub>2</sub>O.

45 30 Mass spectral data was obtained on a Sciex API 365 LC-MS/MS  
system (Toronto, Canada). Separations were done on a Shimadzu HPLC system  
consisting of two LC-10AD pumps and a SCL-10Avp controller (Shimadzu  
Scientific Instruments, Columbia, MD). Analytes were monitored by UV at 214  
50

5 NM with an ABI 785 detector. HPLC separations were achieved using a BDS-  
Hypersil C8 column (150 × 4.6 mm; 120 Å, Keystone Scientific, Inc.,  
Bellefonte, PA). The mobile phase consisted of H<sub>2</sub>O:methanol (98:2 v:v) in  
10 pump A and H<sub>2</sub>O: methanol (2:92 v:v) in pump B. A linear gradient was used  
5 from 5% B to 30% B over 10 minutes and was held at 30% B for 20 minutes  
before returning to initial conditions (mono-carborane synthesis). The di-  
carborane required a longer gradient using the same mobile phases; 5% B to 65%  
15 B over 25 minutes and held at 65% B for 15 minutes before returning to initial  
conditions. The separations were monitored by UV absorption at 214 NM. The  
10 flow was 1.0 ml/minute and was split post-column allowing ~ 10 µl to flow into  
the mass spectrometer.

20 Mass spectral data was collected using electrospray ionization in  
positive mode over a mass range of 300 to 2300 AMU at a dwell time of 0.3  
ms/0.1 AMU. Synthetic samples were prepared at 5 or 10 mg/ml in pump A  
15 mobile phase and an aliquot injected onto the HPLC (1-5 µl). Retention times of  
25 the mono- and di-carborane products were determined to be 13.3 minutes and  
16.2 minutes respectively. Purification of the b and d mono-carborane products  
was achieved by collecting fractions at the elution times for several injections.  
30 The collected fractions were combined and dried to a powder. A portion of the  
20 purified product was dissolved in methanol:H<sub>2</sub>O (1:1) and reanalyzed by HPLC-  
MS to ascertain the purity.

35 α-Carborane, butyllithium (1.6 M solution in hexanes) and  
putrescine were purchased from Aldrich Chemical Company (Milwaukee, WI).  
The water-soluble carbodiimide 1-ethyl-3-(3'-dimethylaminopropyl)  
25 carbodiimide and 1-hydroxybenzotriazole were from Sigma (St. Louis, MO).  
40 Thin layer chromatography (TLC) silica gel plates were obtained from Eastman  
Kodak Company. The cyanocobalamin-b, d, and e monocarboxylic acids and  
the b, d-dicarboxylic acid were prepared as described before (see U.S. Patent  
45 Number 5,739,313, and D.L. Anton, et al., *J. Am. Chem. Soc.*, 1980, 102, 2212-  
30 2219), and were provided by Professor Kahl of the Department of Medicinal  
Chemistry at the University of California, San Francisco.

**Example 1**                      Cyanocobalamin-nido-carborane conjugates (5, Figure 2).

Separate reaction mixtures containing 1 g (~0.66 mmol) of the b-monocarboxylic acid, d-monocarboxylic acid, or the b,d-dicarboxylic acid of cyanocobalamin, hydroxybenzotriazole (810 mg, 6.0 mmol), 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide (11.4 g, 6.0 mmol) and 4 (600 mg, 2.0 mmol) in 100 mL of a water-acetone mixture (2:1) were adjusted to pH 6.9 with 1N NaOH. The reactions were stirred at room temperature and their progress monitored by TLC. After 3 hours, the mixtures were concentrated to remove acetone and the resulting aqueous suspensions were extracted into 92% aqueous phenol. The phenol phases were washed with water to remove the water-soluble reactants. One volume of acetone and three volumes of ether were added to each of the phenol phases and the desired cyanocobalamin-nido-carborane conjugates were back extracted into water. The aqueous layers were extracted three times with ether to remove residual phenol and unreacted (4). Finally the aqueous solutions were evaporated to dryness. The residuals were triturated with acetone and the desired conjugates isolated as orange-red powders (yields 90-95% based on the cyanocobalamin-carboxylic acids).

The isotope ratios of the final products were also compared to theoretical as follows: Mono product theoretical  $M+H^+$   
(C70H109CoB9O15PN15) - 1584.8 (12%), 1585.8 (37%), 1586.8 (74%), 1587.8 (100%), 1588.8 (84%), 1589.8 (44%), 1590.8 (16%), 1591.8 (4%). Synthesis I - 1584.8 (14%), 1585.8 (37%), 1586.8 (74%), 1587.8 (100%), 1588.8 (88%), 1589.8 (48%), 1590.8 (19%), 1591.8 (6%). Synthesis II - 1584.8 (15%), 1585.8 (40%), 1586.8 (78%), 1587.8 (100%), 1588.8 (86%), 1589.8 (44%), 1590.8 (17%), 1591.8 (7%).

Di product theoretical  $M+H^+$  - (C77H129CoB18O16PN16) - 1815.1 (9%), 1816.1 (23%), 1817.1 (47%), 1818.1 (77%), 1819.1 (99%), 1820.1 (100%), 1821.1 (77%), 1822.1 (44%), 1823.1 (19%), 1824.1 (6%). Observed synthesis - 1815.1 (15%), 1816.1 (27%), 1817.1 (51%), 1818.1 (79%), 1819.1 (100%), 1820.1 (100%), 1821.1 (78%), 1822.1 (49%), 1823.1 (25%), 1824.1 (12%).

Results included the identification of the *b* and *d* carborane cyanocobalamin (CCC) analogs by LC-MS. Products identified were

subsequently separated and purified by HPLC. The purified products were reanalyzed by LC-MS with no starting material detected. MS/MS data was also obtained on the *b* and *d* CCC analogs to provide further structural characterization. These purified products were then tested in the biological assays. The *e*-CCC analog and the *di*-CCC analog were also separated and identified by LC-MS, however, the preparations contained more reaction side products resulting in difficult purification. The isotope ratios of the *b*- and *d*-CCC analogs, as well as the *di*-CCC analog, were compared to theoretical isotope ratios to provide further information for identification.

Mass spectra confirmed the presence of each cyanocobalamin-*nido*-carborane conjugate. Mass spectroscopy analysis as well as  $^{11}\text{B}$  NMR showed that during the conversion of 3 to 4 (Figures 2 and 3) the *o*-carborane nucleus lost a boron atom to yield the *nido*-carborane derivative 4 (Figure 3). UV-visible spectroscopy of the final products showed maxima typical of cyanocobalamin, indicating that the corrin nucleus was intact and that the 5,6-dimethyl benzimidazole nucleotide was still attached to the cobalt atom.

The starting material 4 was prepared as follows (see Figure 2):

a. *o*-carborane carboxylic acid (2). A solution of *o*-carborane (1) (5.0 g, 34.7 mmol) in 500 ml dry ether in a 1 L round bottom flask was cooled to  $-78^{\circ}\text{C}$  in a dry ice-acetone bath. The solution was flushed with argon and the flask sealed with a serum stopper. *n*-Butyllithium (24 mL, 1.6 M in hexanes) was slowly injected via a syringe over a period of about 20 minutes and the reaction was stirred for an additional 30 minutes. Crushed dry ice (10-15 g) was then added and the mixture stirred for 1 hour. The dry ice-acetone bath was removed and the reaction allowed to come to room temperature. The ether and hexanes were removed on a rotary evaporator, water (150 ml) was added and unreacted *o*-carborane was removed by extraction with hexanes ( $2 \times 100$  ml). The aqueous phase was acidified with concentrated HCl and the desired product extracted with hexanes ( $4 \times 100$  ml). The combined extracts were dried over  $\text{Na}_2\text{SO}_4$  and evaporated to dryness to give 5.8 g (88.7%) of *o*-carborane carboxylic acid (2), which was used without further purification.

b. o-carborane carboxylic acidchloride (3). o-Carborane carboxylic acid (2.0 g, 10.6 mmol), dried over  $P_2O_5$ , was dissolved in 30 ml thionylchloride and heated under reflux for 3 hours. The solution was cooled to room temperature, evaporated to dryness and dried over  $P_2O_5$  (2.05 g, 9.9 mmol, 93%) to provide (3), which was used without further purification.

c. Nido-Carboranoyl(4-aminobutyl) amide (4). The acid chloride (3) was dissolved in 15 ml dry pyridine and 1,4-diaminobutane (1.12 g, 12.7 mmol) was added. The reaction mixture was heated under reflux for 3 hours, cooled to room temperature and concentrated. Water (10 ml) was added and the suspension acidified with 3 M HCl. The desired product was extracted with ethylacetate (4 × 50 ml), the combined organic layers were washed once with water, dried over  $Na_2SO_4$  and evaporated to dryness to yield 2.35 g (8.0 mmol, 75%) of (4). Thus far the product has resisted crystallization from a variety of solvent mixtures, however TLC on silica gel plates (2-propanol- $NH_4OH-H_2O$ ; 7:1:2) showed only one ninhydrin positive compound distinct from the diamine.

**Example 2. 6-Carboranoylamidopropyl cobalamin.**

6-(3-aminoprop-1-yl)cobalamin (300 mg), hydroxybenzotriazole (270 mg), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (382 mg) and o-carborane monocarboxylic acid were dissolved in water (50 ml) and acetone (20 ml) and allowed to react at room temperature for 3 hours. The reaction mixture was concentrated to remove acetone and desalted via phenol extraction. The concentrated aqueous solution was crystallized from aqueous acetone to give the title compound.

The intermediate 6-(3-aminoprop-1-yl)cobalamin was prepared as follows:

a. 6-(3-aminoprop-1-yl)cobalamin. Cyano cobalamin (500 mg) was dissolved in 100 ml deoxygenated water containing 10 mg  $CoCl_2$ , and reduced with sodium borohydride to give the corresponding Co (I) compound. After 30 minutes, 3-chloropropyl amine hydrochloride (130 mg) dissolved in 5 ml deoxygenated ethanol was added. After one hour, at room temperature, the mixture was desalted via phenol extraction. The aminopropylcarbolamine was back extracted into water after the addition of 1 volume of acetone and 3

5 volumes of ether. The aqueous solution was concentrated and the desired  
6-(3-aminoprop-1-yl)cobalamin crystallized from aqueous acetone (yield 510  
mg).

**Example 3 DTPA-aminopropylcobalamin (DAPC)**

10 5 Using a coupling procedure similar to that described in Example  
2, 6-(3-aminoprop-1-yl)cobalamin and DTPA were coupled in the presence of  
hydroxybenzotriazole to give the title compound.

**Example 4 *In vitro* biological activity of the carborane cyanocobalamin  
analogs.**

10 To assess the *in vitro* binding of the carborane cyanocobalamin  
(Example 1, CCC) and DTPA-aminopropylcobalamin (Example 3, DAPC)  
20 analogs to the transcobalamin proteins, the unsaturated Vitamin B12 binding  
capacity (UBBC) assay (see D.A. Collins and H.P.C. Hogenkamp, *J. Nuclear  
Medicine*, 1997, 38, 717-723) was performed. Serum was obtained from 5  
15 patients being evaluated for pernicious anemia at the Mayo Clinic. The patients'  
serum first underwent a routine clinical UBBC as previously described. To  
25 determine if the CCC and DAPC analogs would inhibit Co-57 cyanocobalamin  
from binding to the transcobalamin proteins, the excess serum from the 5  
patients underwent modified UBBC.

20 Specifically, under dim light, 0.4 ml serum was treated with 4  $\mu$ l  
(concentration 10  $\mu$ g CCC per ml normal saline) of the 2 CCC analogs,  
carborane-d-cyanocobalamin and carborane-b-cyanocobalamin, and DAPC. The  
35 analogs were incubated for 20 minutes at room temperature with the patient's  
serum. Both the clinical run and the analog treated samples were assayed for  
25 UBBC as usual.

40 The analogs competitively blocked Co-57-cyanocobalamin from  
binding to the transcobalamin proteins. Therefore, the cpm of the modified  
UBBC assay was significantly lower than that of the clinical runs. The percent  
PB of the analogs to transcobalamin protein was calculated as follows ( $PB = 100$   
45  $- CCC_{UBBC} \text{ cpm} / \text{clinical UBBC cpm} \times 100$ ). The average percent binding (PB)  
30 of the 5 solutions ( $n = 10$  for each solution, i.e., two modified UBBC assays per  
patient) for the carborane-d cyanocobalamin and carborane-b cyanocobalamin  
was 35.75% and 92.93% respectively, and 98.21% for DAPC.



5 All publications, patents, and patent documents are incorporated  
by reference herein, as though individually incorporated by reference. The  
invention has been described with reference to various specific and preferred  
embodiments and techniques. However, it should be understood that many  
10 5 variations and modifications may be made while remaining within the spirit and  
scope of the invention.

## Claims

5

10

15

20

25

30

35

40

45

50

55

What is claimed is:

1. A compound wherein a residue of a compound of formula I (Figure 1) is linked to a residue of a molecule comprising B-10, wherein X is CN, OH, CH<sub>3</sub>, adenosyl, or a molecule comprising B-10; or a pharmaceutically acceptable salt thereof.
2. The compound of claim 1 wherein the residue of a molecule comprising B-10 is directly linked to the 6-position of the compound of formula I or is directly linked to a residue of the b, d or e-carboxamide group of the compound of formula I.
3. The compound of claim 1 wherein the residue of a molecule comprising B-10 is linked by a linker to the 6-position of the compound of formula I or is linked by a linker to the residue of a-, b-, d- or e-carboxamide group of the compound of formula I.
4. The compound of claim 1 wherein the residue of a molecule comprising B-10 is linked to a residue of the b-carboxamide group of the compound of formula I.
5. The compound of claim 1 wherein the residue of a molecule comprising B-10 is linked to a residue of d-carboxamide group of the compound of formula I.
6. The compound of claim 1 wherein the residue of a molecule comprising B-10 is linked to a residue of the e-carboxamide group of the compound of formula I.
7. The compound of claim 1 wherein a residue of a molecule comprising B-10 is linked to a residue of the b-carboxamide group and a second residue of a molecule comprising B-10 is linked to a residue of the d-carboxamide group of the compound of formula I.

- 5 8. The compound of claim 1 wherein the residue of a molecule comprising B-10 is linked to the 6-position of the compound of formula I.
- 10 9. The compound of claim 1 wherein the molecule comprising B-10 contains 1 to about 20 boron atoms, inclusive.
- 15 10. The compound of claim 1 wherein the molecule comprising B-10 is an amino acid, a carbohydrate, a nucleoside or a carborane.
- 20 11. The compound of claim 1 wherein the molecule comprising B-10 is o-carborane, m-carborane or p-carborane.
- 25 12. The compound of claim 1 wherein the molecule comprising B-10 is o-carborane.
- 30 13. The compound of claim 3 wherein at least one linker is of the formula W-A-Q wherein A is (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>2</sub>-C<sub>6</sub>)alkynyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, or (C<sub>6</sub>-C<sub>10</sub>)aryl, wherein W and Q are each independently -N(R)C(=O)-, -C(=O)N(R)-, -OC(=O)-, -C(=O)O-, -O-, -S-, -S(O)-, -S(O)<sub>2</sub>-, -N(R)-, -C(=O)-, or a direct bond; wherein each R is independently H or (C<sub>1</sub>-C<sub>6</sub>)alkyl.
- 35 14. The compound of claim 13 wherein W is NH<sub>2</sub> or COOH and Q is NH<sub>2</sub> or COOH.
- 40 15. The compound of claim 13 wherein A is (C<sub>1</sub>-C<sub>6</sub>)alkyl.
- 45 16. The compound of claim 3 wherein at least one linker is about 5 angstroms to about 50 angstroms, inclusive.
- 50 17. The compound of claim 3 wherein at least one linker comprises a therapeutic radionuclide or a diagnostic radionuclide.

18. The compound of claim 17 wherein the therapeutic radionuclide is a metallic radionuclide.

19. The compound of claim 17 wherein the diagnostic radionuclide is a metallic radionuclide.

20. The compound of claim 17 wherein the diagnostic radionuclide is a non-metallic radionuclide.

21. The compound of claim 3 wherein at least one linker is a divalent radical formed from a peptide.

22. The compound of claim 3 wherein at least one linker is a divalent radical formed from an amino acid.

23. The compound of claim 3 wherein at least one linker is poly-L-glutamic acid, poly-L-aspartic acid, poly-L-histidine, poly-L-ornithine, poly-L-serine, poly-L-threonine, poly-L-tyrosine, poly-L-lysine-L-phenylalanine, poly-L-lysine or poly-L-lysine-L-tyrosine.

24. The compound of claim 1 wherein the residue of the compound of formula I is also linked to a linker comprising a detectable radionuclide or a therapeutic radionuclide.

25. A compound wherein a residue of a compound of formula I (Figure 1) is linked to a group of the formula Q-L-W-Det; wherein X is CN, OH, CH<sub>3</sub>, adenosyl, a molecule comprising B-10, or Q-L-W-Det; wherein Det is a chelating group comprising Gd-157; L is a linker or absent; and W and Q are each independently -N(R)C(=O)-, -C(=O)N(R)-, -OC(=O)-, -C(=O)O-, -O-, -S-, -S(O)-, -S(O)<sub>2</sub>-, -C(=O)-, -N(R)-, or a direct bond; wherein each R is independently H or (C<sub>1</sub>-C<sub>6</sub>)alkyl; or a pharmaceutically acceptable salt thereof.

26. The compound of claim 25 wherein the group of the formula Q-L-W-Det is linked to a residue of the b-carboxamide group, a residue of the d-carboxamide group, a residue of the e-carboxamide, or the 6-position of the compound of formula I.

27. The compound of claim 25 wherein the group of the formula Q-L-W-Det is linked to a residue of the b-carboxamide group and a second group of the formula Q-L-W-Det is linked to a residue of the d-carboxamide group of the compound of formula I.

28. The compound of claim 25 wherein the group of the formula Q-L-W-Det is between about 20 and about 500 angstroms, inclusive, in length.

29. The compound of claim 25 wherein at least one chelating group is EDTA, DTPA, DOTA, DOTMP, TETA, MAG3 or DCTA.

30. The compound of claim 25 wherein at least one chelating group is DTPA comprising Gd-157.

31. A compound wherein a residue of a compound of formula I (Figure 1) is linked to a residue of a molecule comprising B-10; wherein a residue of the compound of formula I is also linked to a group of the formula Q-L-W-Det, wherein X is CN, OH, CH<sub>3</sub>, adenosyl, a group of the formula Q-L-W-Det, or a molecule comprising B-10; wherein:

Det is a chelating group comprising a therapeutic radionuclide or a diagnostic radionuclide;

L is a linker or absent; and

Q and W are each independently -N(R)C(=O)-, -C(=O)N(R)-, -OC(=O)-, -C(=O)O-, -O-, -S-, -S(O)-, -S(O)<sub>2</sub>-, -C(=O)-, -N(R)-, or a direct bond; wherein each R is independently H or (C<sub>1</sub>-C<sub>6</sub>)alkyl; or a pharmaceutically acceptable salt thereof.

5 32. The compound of claim 31 wherein at least one of the radionuclides is Tc<sup>99m</sup>, In<sup>111</sup>, In<sup>110</sup>, Gd<sup>157</sup> or Y<sup>86</sup>.

10 33. The compound of claim 31 wherein a molecule comprising B-10 is linked to the residue of a b-carboxamide group, d-carboxamide group, e-carboxamide group, or the 6-position of the compound of formula I.

15 34. The compound of claim 31 wherein the chelating group is DTA, DTPA, DOTA, TETA, DOTMP, MAG3 or DCTA.

20 35. The compound of claim 31 wherein the chelating group is DTPA.

25 36. The compound of claim 31 wherein the residue of a molecule comprising B-10 contains 1 to about 20 boron atoms.

30 37. The compound of claim 31 wherein the molecule comprising B-10 is a carbohydrate, an amino acid, a nucleoside or a carborane.

35 38. The compound of claim 31 wherein the molecule comprising B-10 is o-nido-carborane, m-nido-carborane or p-nido-carborane.

40 39. The compound of claim 31 wherein the molecule comprising B-10 is o-carborane.

45 40. The compound of claim 31 wherein the residue of a molecule comprising B-10 is directly linked to the 6-position or to the residue of the b-, d- or e-carboxamide group of the compound of formula I.

50 41. The compound of claim 31 wherein the residue of the compound of formula I is linked to a residue of a molecule comprising B-10 through a linker.

5 42. The compound of claim 41 wherein the linker comprises a non-metallic radionuclide.

10 43. The compound of claim 41 wherein the linker is about 5 angstroms to about 50 angstroms, inclusive.

15 44. The compound of claim 1 further comprising a detectable radionuclide.

20 45. The compound of claim 2 wherein the detectable radionuclide is a non-metallic radionuclide.

25 46. The compound of claim 3 wherein the non-metallic radionuclide is Carbon-11, Fluorine-18, Bromine-76, Iodine-123, or Iodine-124.

47. The compound of claim 2 wherein the detectable radionuclide is directly linked to the compound of formula I.

30 48. The compound of claim 2 wherein the detectable radionuclide is linked by a linker to the compound of formula I.

35 49. The compound of claim 6 wherein the linker is of the formula W-A wherein A is (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>2</sub>-C<sub>6</sub>)alkynyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, or (C<sub>6</sub>-C<sub>10</sub>)aryl, wherein W is -N(R)C(=O)-, -C(=O)N(R)-, -OC(=O)-, -C(=O)O-,  
25 -O-, -S-, -S(O)-, -S(O)<sub>2</sub>-, -N(R)-, -C(=O)-, or a direct bond; wherein each R is independently H or (C<sub>1</sub>-C<sub>6</sub>)alkyl; and wherein A is substituted with one or more  
40 non-metallic radionuclides.

45 50. The compound of claim 48 wherein the linker is about 5 angstroms to about 50 angstroms, inclusive.

50 51. The compound of claim 48 wherein the linker is a divalent peptide or amino acid.



5 52. The compound of claim 48 wherein the linker is poly-L-glutamic acid, poly-L-aspartic acid, poly-L-histidine, poly-L-ornithine, poly-L-serine, poly-L-threonine, poly-L-tyrosine, poly-L-lysine-L-phenylalanine, or poly-L-lysine or poly-L-lysine-L-tyrosine.

10 53. The compound of claim 48 wherein the linker is linked to the 6-position of the compound of formula I or is linked to the residue of a-, b-, d- or e-carboxamide group of the compound of formula I.

15 54. A compound wherein a residue of a compound of formula I (Figure 1) is linked 1) to a detectable radionuclide; and is linked 2) to a group comprising Gd-157; or a pharmaceutically acceptable salt thereof.

20 55. The compound of claim 54 wherein the group comprising Gd-157 has the formula Q-L-W-Det; wherein X is CN, OH, CH<sub>3</sub>, adenosyl, a molecule comprising B-10, or Q-L-W-Det; Det is a chelating group comprising Gd-157; L is a linker or absent; and W and Q are each independently -N(R)C(=O)-, -C(=O)N(R)-, -OC(=O)-, -C(=O)O-, -O-, -S-, -S(O)-, -S(O)<sub>2</sub>-, -C(=O)-, -N(R)-, or a direct bond; wherein each R is independently H or (C<sub>1</sub>-C<sub>6</sub>)alkyl.

25 56. The compound of claim 54 wherein the detectable radionuclide is a non-metallic radionuclide.

30 57. The compound of claim 56 wherein the non-metallic radionuclide is Carbon-11, Fluorine-18, Bromine-76, Iodine-123, or Iodine-124.

35 58. The compound of claim 54 wherein the detectable radionuclide is directly linked to the compound of formula I.

40 59. The compound of claim 54 wherein the detectable radionuclide is linked by a linker to the compound of formula I.

5 60. The compound of claim 59 wherein the linker is of the formula  
W-A wherein A is (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>2</sub>-C<sub>6</sub>)alkynyl, (C<sub>3</sub>-  
C<sub>3</sub>)cycloalkyl, or (C<sub>6</sub>-C<sub>10</sub>)aryl, wherein W is -N(R)C(=O)-, -C(=O)N(R)-,  
-OC(=O)-, -C(=O)O-, -O-, -S-, -S(O)-, -S(O)<sub>2</sub>-, -N(R)-, -C(=O)-, or a direct  
10 5 bond; wherein each R is independently H or (C<sub>1</sub>-C<sub>6</sub>)alkyl; and wherein A is  
substituted with One or more non-metallic radionuclides.

15 61. The compound of claim 59 wherein the linker is about 5  
angstroms to about 50 angstroms, inclusive.

10 62. The compound of claim 59 wherein the linker is a divalent  
peptide or amino acid.

25 63. The compound of claim 59 wherein the linker is poly-L-glutamic  
acid, poly-L-aspartic acid, poly-L-histidine, poly-L-ornithine, poly-L-serine,  
poly-L-threonine, poly-L-tyrosine, poly-L-lysine-L-phenylalanine, or poly-L-  
lysine or poly-L-lysine-L-tyrosine.

30 64. The compound of claim 59 wherein the linker is linked to the 6-  
20 position of the compound of formula I or is linked to the residue of a-, b-, d- or  
e-carboxamide group of the compound of formula I.

35 65. A compound wherein a residue of a compound of formula I  
(Figure 1) is linked 1) to a molecule comprising B-10 or to a chelating group  
25 comprising Gd-157, and 2) to at least one residue of the formula Q-L-W-Det;  
wherein each Det is independently a chelating group comprising a metallic  
40 radionucleotide; each L is independently a linker or absent; and each W and Q is  
independently -N(R)C(=O)-, -C(=O)N(R)-, -OC(=O)-, -C(=O)O-, -O-, -S-,  
-S(O)-, -S(O)<sub>2</sub>-, -C(=O)-, -N(R)-, or a direct bond; wherein each R is  
45 30 independently H or (C<sub>1</sub>-C<sub>6</sub>)alkyl; or a pharmaceutically acceptable salt thereof.

50 66. The compound of claim 1 or 44 wherein the residue of a compound of  
formula I is also linked to a group comprising Gd-157.

5 67. The compound of claim 66 wherein the group comprising Gd-157  
has the formula Q-L-W-Det; wherein X is CN, OH, CH<sub>3</sub>, adenosyl, a molecule  
comprising B-10, or Q-L-W-Det; Det is a chelating group comprising Gd-157; L  
is a linker or absent; and W and Q are each independently -N(R)C(=O)-,  
10 5 -C(=O)N(R)-, -OC(=O)-, -C(=O)O-, -O-, -S-, -S(O)-, -S(O)<sub>2</sub>-, -C(=O)-, -N(R)-,  
or a direct bond; wherein each R is independently H or (C<sub>1</sub>-C<sub>6</sub>)alkyl.

15 68. A pharmaceutical composition comprising a compound of any  
one of claims 1-67 and a pharmaceutically acceptable carrier.

10 69. A method of treating a tumor in a mammal in need of such  
treatment comprising administering to the mammal an effective amount of a  
20 compound of any one of claims 1-67 in combination with a pharmaceutically  
acceptable vehicle; and administering neutron capture therapy.

15 70. A method for imaging a tumor in a mammal comprising  
administering to the mammal a detectable amount of a compound of any one of  
25 claims 1-67; and detecting the presence of the compound.

30 71. The method of claim 70 further comprising treating the tumor  
with neutron capture therapy.

35 72. A compound of any one of claims 1-67 for use in medical therapy  
or diagnosis.

25 73. The use of a compound of any one of claims 1-67 for the  
40 manufacture of a medicament for imaging a tumor in a mammal.

45 74. The use of a compound of any one of claims 1-67 for the  
30 manufacture of a medicament for treating a tumor in a mammal in need of such  
treatment.





FIG. 1

2/3



(o-CARB) = o-Carborane

[o-nido-CARB]<sup>-</sup> = o-nido-Carborane

FIG. 2




FIG. 3